arXiv:0709.1262v1 [math.AG] 9 Sep 2007

An $O(2,n)$ formulation of invariant theory for elliptic Weyl group

Ikuo Satake

Department of Mathematics, Osaka University, Toyonaka, Osaka,
560-0043, JAPAN

Abstract

In this paper, we give a new formulation of invariant theory for elliptic Weyl group using the group $O(2,n)$. As an elliptic Weyl group quotient, we define a suitable $\mathbb{C}^*$-bundle. We show that it has a conformal Frobenius structure which we define in this paper. Then its good section could be identified with a Frobenius manifold which we constructed in [13].

1 Introduction

Motivated by a period mapping for a primitive form [9] for an unfolding of a function with a simple elliptic singularity, an elliptic root system is introduced [10]. On the quotient space $\tilde{\mathbb{E}}_a//W$ of the domain $\tilde{\mathbb{E}}_a$ by an elliptic Weyl group W, a flat holomorphic metric is constructed by [11] and Frobenius manifold structure is constructed by [13].

Here the domain $\tilde{\mathbb{E}}_a$ is not canonical, that is, it depends on the choice of a marking a. Then we set the following problems.

Problem 1: What is a relation between the elliptic Weyl group quotient spaces $\tilde{\mathbb{E}}_a//W$ and $\tilde{\mathbb{E}}_{a'}//W^2$ for different markings a and a'?

Problem 2: What is a relation between the Frobenius manifold structure on them?

By definition, an elliptic root system is a root system belonging to a vector space with an inner produce with a signature $(0,2,l)$ where we denote by (l_+,l_0,l_-) numbers of positive, 0 and negative eigenvalues respectively (we choose a negative semi-definite inner product). A domain $\tilde{\mathbb{E}}_a$ corresponds to a hyperbolic extension with a signature $(1,1,l+1)$.

In this paper, we consider 2-extension with a signature $(2,0,l+2)$ and construct a $\mathbb{C}^*$-bundle

$$p : D^2 \to \mathbb{P}(D^2), \tag{1.1}$$

where D^2 is a quadric hypersurface of a suitable domain D and $\mathbb{P}(D^2) = D^2/\mathbb{C}^*$. They corresponds to the group $O(2,0,l+2)$. Then we find that $\tilde{\mathbb{E}}_a$ could be canonically identified with a section of p.

This formulation is good from the viewpoint of the group action of the automorphism of an elliptic root system. We find that a central extension group of the group of the automorphisms of an elliptic root system is naturally defined in our formulation and this group acts on the $\mathbb{C}^*$-bundle p equivariantly. The fact that this action changes a section of p, corresponds to the fact that this group is not contained in $O(1,1,l+1)$.

We see a Frobenius manifold structure. The elliptic Weyl group is constructed as a subgroup of $O(2,0,n)$ and denote it by W^2, which is isomorphic to W. Then the Weyl group quotient:

$$p : D^2//W^2 \to \mathbb{P}(D^2)//W^2 \tag{1.2}$$

is again a $\mathbb{C}^*$-bundle and the space $\tilde{\mathbb{E}}_a//W$ gives a section of it. Since $\tilde{\mathbb{E}}_a//W$ is a Frobenius manifold, the base space $\mathbb{P}(D^2)//W^2$ is also a Frobenius manifold. But if we choose another marking a' and a corresponding another section $\tilde{\mathbb{E}}_{a'}//W$, then $\mathbb{P}(D^2)//W^2$ has a structure of another Frobenius manifold. In fact, except holomorphic metrics, these two Frobenius manifold structures are same. For holomorphic metrics, they are conformally equivalent.

Thus we introduce a notion of conformal Frobenius structure and we prove that $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ has essentially unique conformal Frobenius structure. For any marking a, a Frobenius manifold structure of $\tilde{\mathbb{E}}_a//W$ is obtained from this conformal Frobenius manifold structure. This clarifies the relation of the Frobenius manifold structures of $\tilde{\mathbb{E}}_a//W$ and $\tilde{\mathbb{E}}_{a'}//W^2$.

Also we see that the conformal Frobenius structure has an action of a (central extension of) automorphism group of the elliptic root system. This clarifies the meaning of a conformal transformation in [12] (see Remark 5.4).

This paper is organaized as follows.

In Section 2, we define 2-extension for elliptic root system and define an elliptic Weyl group W^2 and a $\mathbb{C}^*$-bundle $p : D^2 \to \mathbb{P}(D^2)$. We introduce open subsets of a $\mathbb{C}^*$-bundle $p : \overset{\circ}{D^2} \to \mathbb{P}(\overset{\circ}{D^2})$ whose complements are zero of roots. Then we take an elliptic Weyl group quotient $p : \overset{\circ}{D^2} /W^2 \to \mathbb{P}(\overset{\circ}{D^2})/W^2$. We call them "analytic Weyl group quotient spaces". On the space $\overset{\circ}{D^2} /W^2$, we define a tensor $I^*_{\overset{\circ}{D^2}/W^2}$ derived from an inner product of the elliptic root system.

In Section 3, we define elliptic Weyl group invariant rings. We define "algebraic Weyl group quotient spaces" $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ by taking Specan of these elliptic Weyl group invariant rings. On the space $D^2//W^2$, we also define a tensor $I^*_{D^2//W^2}$. derived from an inner product of the elliptic root system. Our theorems are stated for "algebraic Weyl group quotient spaces". But for proofs, we utilize "analytic Weyl group quotient spaces".

In Section 4, we introduce a notion of conformal Frobenius structure and state that a $\mathbb{C}^*$-bundle $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ has essentially unique conformal Frobenius structure. We also show that a group of central extension of automorphism group of the elliptic root system acts on the conformal Frobenius structure.

In Section 5, We give proofs. We relate our formulation with a formulation of [11] and [13] in which the space $\tilde{\mathbb{E}}_a$ is used. We obtain the existence of a conformal Frobenius structure by this comparison. For a uniqueness theorem, we use a conformal deformation of a holomorphic metric of a Frobenius manifold.

We use the following terminologies.

For a complex manifold M, we denote by $\mathcal{O}_M$ (resp. Ω^1_M, Θ_M) the sheaf of holomorphic functions (resp. holomorphic 1-forms, holomorphic vector fields). We denote by $\mathcal{O}(M)$ (resp. $\Omega^1(M)$, $\Theta(M)$) the module $\Gamma(M, \mathcal{O}_M)$ (resp. $\Gamma(M, \Omega^1_M)$, $\Gamma(M, \Theta_M)$).

Let $\mathcal{F}$ be a sheaf on the complex manifold N. For a morphism $f : M \to N$ of complex manifolds M and N, we denote by f^* the following natural morphism

$$f^* : \Gamma(N, \mathcal{F}) \to \Gamma(M, f^*\mathcal{F}). \tag{1.3}$$

induced by $\mathcal{F} \to f_* f^* \mathcal{F}$.

For an isomorphism $f : M \to M$, we have canonical isomorphisms $f^*\Omega^1_M \xrightarrow{\sim} \Omega^1_M$ and $f^*\Theta_M \xleftarrow{\sim} \Theta_M$. Then the morphism (1.3) induces

$$f^* : \Gamma(M, (\Omega^1_M)^{\otimes p} \otimes_{\mathcal{O}_M} (\Theta_M)^{\otimes q}) \xrightarrow{\sim} \Gamma(M, (\Omega^1_M)^{\otimes p} \otimes_{\mathcal{O}_M} (\Theta_M)^{\otimes q}). \tag{1.4}$$

For a $\mathbb{C}^*$-bundle $p : L \to M$ and the isomorphism f of the $\mathbb{C}^*$-bundle:

$$\begin{tikzcd} L \arrow[r, "f", "\sim"'] \arrow[d, "p"'] & L \arrow[d, "p"'] \\ M \arrow[r, "f", "\sim"'] & M \end{tikzcd} , \tag{1.5}$$

we have $f^*(p^*\Omega^1_M) \simeq p^* f^* \Omega^1_M \xrightarrow{\sim} p^*\Omega^1_M$ and $f^*(p^*\Theta_M) \simeq p^* f^* \Theta_M \xleftarrow{\sim} p^*\Theta_M$. Then the morphism (1.3) induces

$$f^* : \Gamma(L, (p^*\Omega^1_M)^{\otimes p} \otimes_{\mathcal{O}_L} (p^*\Theta_M)^{\otimes q}) \xrightarrow{\sim} \Gamma(L, (p^*\Omega^1_M)^{\otimes p} \otimes_{\mathcal{O}_L} (p^*\Theta_M)^{\otimes q}). \tag{1.6}$$

2 Analytic Weyl group quotient space

2.1 Elliptic root system

In this subsection, we define an elliptic root system, its orientation and 2-extension.

2.1.1 Definition of elliptic root system

Let l be a positive integer. Let F be a real vector space of rank $l+2$ with a negative semi-definite or positive semi-definite symmetric bilinear form $I_F : F \times F \to \mathbb{R}$, whose radical $\mathrm{rad} I_F := \{x \in F \mid I_F(x, y) = 0, \forall y \in F\}$ is a vector space of rank 2. For a non-isotropic element $\alpha \in F$ (i.e. $I_F(\alpha, \alpha) \neq 0$), we put $\alpha^\vee := 2\alpha / I_F(\alpha, \alpha) \in F$. The reflection w_α with respect to α is defined by

$$w_\alpha(u) := u - I_F(u, \alpha^\vee)\alpha \quad (\forall u \in F). \tag{2.1}$$

Definition 2.1. ([10, p.104, Def. 1]) *A set R of non-isotropic elements of F is an elliptic root system belonging to (F, I_F) if it satisfies the axioms 1-4:*

1. *The additive group generated by R in F, denoted by $Q(R)$, is a full sublattice of F. That is, the embedding $Q(R) \subset F$ induces the isomorphism : $Q(R) \otimes_{\mathbb{Z}} \mathbb{R} \simeq F$.*
2. *$I_F(\alpha, \beta^\vee) \in \mathbb{Z}$ for $\alpha, \beta \in R$.*
3. *$w_\alpha(R) = R$ for $\forall \alpha \in R$.*
4. *If $R = R_1 \cup R_2$, with $R_1 \perp R_2$, then either R_1 or R_2 is void.*

2.1.2 Definition of orientation

For an elliptic root system R belonging to (F, I_F), the additive group $\mathrm{rad} I_F \cap Q(R)$ is isomorphic to $\mathbb{Z}^2$.

Definition 2.2. *An elliptic root system R is called oriented if the $\mathbb{R}$-vector space $\mathrm{rad} I_F$ is oriented. An ordered pair $\{a, b\}$ of $\mathrm{rad} I_F$ is called an oriented basis if it gives an $\mathbb{R}$-basis of $\mathrm{rad} I_F$ and it gives the orientation of $\mathrm{rad} I_F$.*

2.1.3 Definition of a signed marking

Definition 2.3. *Let R be an elliptic root system R belonging to (F, I_F). By a signed marking, we mean a non-zero element a of $\mathrm{rad} I_F \cap Q(R)$ such that $Q(R) \cap \mathbb{R}a = \mathbb{Z}a$ and the quotient root system $R/\mathbb{R}a$ ($:= \mathrm{Image}(R \hookrightarrow F \to F/\mathbb{R}a)$) is reduced (i.e. $\alpha, c\alpha \in R/\mathbb{R}a$ implies $c \in \{\pm 1\}$). We denote the set of a signed marking by $\Lambda_{\mathbb{Z}}$.*

Hereafter we assume that $\Lambda_{\mathbb{Z}} \neq \emptyset$.

2.1.4 2-extension

We define a 2-extension. Let F^2 be a real vector space of rank $l + 4$ and $I_{F^2} : F^2 \times F^2 \to \mathbb{R}$ an $\mathbb{R}$-symmetric bilinear form. The pair (F^2, I_{F^2}) is called a 2-extension of (F, I_F) if F^2 contains F as a linear subspace, $\mathrm{rad} I_{F^2} = \{0\}$ and $I_{F^2}|_F = I_F$. I_{F^2} is indefinite symmetric bilinear form with signature of $(2, l+2)$ or $(l+2, 2)$ according to I is negative semi-definite or positive semi-definite. A 2-extension is unique up to isomorphism. Hereafter we fix a hyperbolic extension (F^2, I_{F^2}).

2.2 Orthogonal group and Weyl group

2.2.1 Definition of an Orthogonal group and a Weyl group

In this subsection, we define a parabolic subgroup of an orthogonal group and Weyl group.

Let $F^2_{\mathbb{C}} := F^2 \otimes_{\mathbb{R}} \mathbb{C}$, $(\mathrm{rad} I_F)_{\mathbb{C}} := \mathrm{rad} I_F \otimes_{\mathbb{R}} \mathbb{C}$. $I_{F^2_{\mathbb{C}}}$ is a $\mathbb{C}$-bilinear extension of I_{F^2}. We put $\tilde{w}_\alpha(u) := u - I_{F^2_{\mathbb{C}}}(u, \alpha^\vee)\alpha$ for $u \in F^2_{\mathbb{C}}$.

We define the groups

$$SL(\mathrm{rad} I_F) := \{ g \in GL(\mathrm{rad} I_F) \mid \det g = 1 \}, \tag{2.2}$$

$$O(F) := \{ g \in GL(F) \mid I_F(gx, gy) = I_F(x, y), \forall x, y \in F \}, \tag{2.3}$$

$$O^+(F) := \{ g \in O(F) \mid g|_{\mathrm{rad} I_F} \in SL(\mathrm{rad} I_F) \}, \tag{2.4}$$

$$Aut(R) := \{ g \in GL(F) \mid g(R) = R,\ g \in O^+(F) \}, \tag{2.5}$$

$$O(F^2_{\mathbb{C}}) := \{ g \in GL(F^2_{\mathbb{C}}) \mid I_{F^2_{\mathbb{C}}}(gx, gy) = I_{F^2_{\mathbb{C}}}(x, y),\ \forall x, y \in F^2_{\mathbb{C}} \}, \tag{2.6}$$

$$O^+(F^2_{\mathbb{C}}, F) := \{ g \in O(F^2_{\mathbb{C}}) \mid g(F) \subset F,\ g|_F \in O^+(F) \}, \tag{2.7}$$

$$W^2 := \langle \tilde{w}_\alpha \in O^+(F^2_{\mathbb{C}}, F) \mid \alpha \in R \rangle. \tag{2.8}$$

The natural morphism

$$\pi^2 : O^+(F^2_{\mathbb{C}}, F) \to O^+(F) \tag{2.9}$$

is surjective. We put

$$K^2_{\mathbb{C}} := \ker \pi^2, \tag{2.10}$$

$$K^2_{\mathbb{Z}} := \ker(\pi^2|_{W^2}), \tag{2.11}$$

$$\widetilde{Aut}(R) := (\pi^2)^{-1}(Aut(R)). \tag{2.12}$$

2.2.2 Properties of an orthogonal group

Proposition 2.4. *1) The group $O^+(F^2_{\mathbb{C}}, F)$ acts on the flag $F^2_{\mathbb{C}} \supset F \supset \mathrm{rad} I_F$.*
2) The group $K^2_{\mathbb{C}}$ is isomorphic to additive group $\mathbb{C}$ and it is contained in the center of $O^+(F^2_{\mathbb{C}}, F)$.
3) The group $\widetilde{Aut}(R)$ contains W^2 as a normal subgroup.

Proof. By definition, we obtain 1). We obtain 2) by direct calculation. Since $g\tilde{w}_\alpha g^{-1} = \tilde{w}_{g(\alpha)}$ for $g \in \widetilde{Aut}(R)$, we obtain 3). $\square$

2.3 Description of a center

In this subsection, we construct the canonical group isomorphism $\rho : \mathbb{C} \to K_{\mathbb{C}}$ by the aid of the Eichler-Siegel transformation ([10, p93]).

We define the Eichler-Siegel transformation by

$$ES : F^2_{\mathbb{C}} \otimes_{\mathbb{C}} F^2_{\mathbb{C}} \to \mathrm{End}_{\mathbb{C}}(F^2_{\mathbb{C}}), \quad \sum_i \alpha_i \otimes \beta_i \mapsto u - \sum_i \alpha_i I_{F^2_{\mathbb{C}}}(u, \beta_i). \tag{2.13}$$

Here $F^2_{\mathbb{C}} \otimes_{\mathbb{C}} F^2_{\mathbb{C}}$ has the semi-group structure by the product

$$(\sum_i u_i \otimes v_i) \circ (\sum_j w_j \otimes x_j) = (\sum_i u_i \otimes v_i) + (\sum_j w_j \otimes x_j) - (\sum_{i,j} I_{F^2_{\mathbb{C}}}(v_i, w_j) u_i \otimes x_j). \tag{2.14}$$

Then ES is a semi-group isomorphism.

Proposition 2.5. *1) ES induces the isomorphism $\wedge^2(\mathrm{rad} I_F)_{\mathbb{C}}$ to $K^2_{\mathbb{C}}$.*
2) A semigroup

$$\mathbb{R}_{>0}(\mathrm{sgn}(I_F))(a \otimes b - b \otimes a) \cap ES^{-1}(W^2)\,(\simeq \mathbb{Z}_{>0}) \tag{2.15}$$

has the unique generator g_0, where $\mathrm{sgn}(I_F)$ is ± 1 according to I_F is positive semi-definite or negative semi-definite.
3) The morphism ES and g_0 depend on the normalization of I_F. But the morphism

$$\rho : \mathbb{C} \to K_{\mathbb{C}}, \quad \alpha \mapsto ES(g_0 \otimes \alpha) \tag{2.16}$$

does not depend on a scalar multiplication ($\mathbb{R}^\times$) of I_{F^2}.

Proof. We could interprete the morphism $\pi^2 : O^+(F^2_{\mathbb{C}}, F) \to O^+(F)$ into the spaces $F^2_{\mathbb{C}} \otimes_{\mathbb{C}} F^2_{\mathbb{C}}$ and $F \otimes_{\mathbb{R}} F/\mathrm{rad} I_F$ by the morphism ES. Then we have a result. $\square$

Remark 2.6. *The element g_0 is exactly written by the notion of marked extended affine root system as $g_0 = (I_R : I)\frac{l_{max}+1}{m_{max}}(a \otimes b - b \otimes a)$, if a, b are oriented basis and $\mathrm{rad} I_F \cap Q(R) \simeq \mathbb{Z}a \oplus \mathbb{Z}b$. See [11, p27 (2.6.2)].*

2.4 $\mathbb{C}^*$-bundles

2.4.1 Preparations

We arbitrary fix an oriented basis a, b. We put $(F^2_{\mathbb{C}})^* := \mathrm{Hom}_{\mathbb{C}}(F^2_{\mathbb{C}}, \mathbb{C})$, $(\mathrm{rad} I_F)^*_{\mathbb{C}} := \mathrm{Hom}_{\mathbb{C}}((\mathrm{rad} I_F)_{\mathbb{C}}, \mathbb{C})$. We denote a natural pairing $F^2_{\mathbb{C}} \times (F^2_{\mathbb{C}})^* \to \mathbb{C}$ by $\langle\ ,\ \rangle$. A non-degenerate $\mathbb{C}$-blinear form $I_{(F^2_{\mathbb{C}})}$ gives a dual $\mathbb{C}$-blinear form $I_{(F^2_{\mathbb{C}})^*} : (F^2_{\mathbb{C}})^* \times (F^2_{\mathbb{C}})^* \to \mathbb{C}$.

The group $GL(F^2_{\mathbb{C}})$ gives the left action on the space $(F^2_{\mathbb{C}})^*$ by

$$\langle g^{-1} \cdot \alpha, x \rangle = \langle \alpha, g \cdot x \rangle \tag{2.17}$$

for $\forall \alpha \in F^2_{\mathbb{C}}, \forall x \in (F^2_{\mathbb{C}})^*, \forall g \in GL(F^2_{\mathbb{C}})$.

2.4.2 Definition of $\mathbb{C}^*$ bundles

We put

$$D := \{ x \in (F^2_{\mathbb{C}})^* \mid \langle a, x \rangle \neq 0,\ \langle b, x \rangle \neq 0,\ \mathrm{Im}\frac{\langle b, x \rangle}{\langle a, x \rangle} > 0 \}, \tag{2.18}$$

$$D^2 := \{ x \in D \mid I_{(F^2_{\mathbb{C}})^*}(x, x) = 0 \}, \tag{2.19}$$

$$\mathbb{H}_{half} := \{ x \in (\mathrm{rad} I_F)^*_{\mathbb{C}} \mid \langle a, x \rangle \neq 0,\ \langle b, x \rangle \neq 0,\ \mathrm{Im}\frac{\langle b, x \rangle}{\langle a, x \rangle} > 0 \}. \tag{2.20}$$

These are complex manifolds and do not depend on the choice of an oriented basis a, b, but depend only on the orientation.

We introduce group actions. Since the spaces $(F^2_{\mathbb{C}})^*$ and $(\mathrm{rad} I_F)^*_{\mathbb{C}}$ are $\mathbb{C}$-vector spaces, they have a natural $\mathbb{C}^*$-action. It induces $\mathbb{C}^*$-actions on D^2 and $\mathbb{H}_{half}$:

$$\psi(\alpha) : D^2 \to D^2, \quad x \mapsto \alpha \cdot x \ (\forall \alpha \in \mathbb{C}^*), \tag{2.21}$$

$$\psi(\alpha) : \mathbb{H}_{half} \to \mathbb{H}_{half}, \quad x \mapsto \alpha \cdot x \ (\forall \alpha \in \mathbb{C}^*). \tag{2.22}$$

The group $\widetilde{Aut}(R)$ acts on $(F^2_{\mathbb{C}})^*$ and $(\mathrm{rad} I_F)^*_{\mathbb{C}}$ by (2.17). We could easily check that this action induces the actions

$$\varphi(g) : D^2 \to D^2 \quad (\forall g \in \widetilde{Aut}(R)), \tag{2.23}$$

$$\varphi(g) : \mathbb{H}_{half} \to \mathbb{H}_{half} \quad (\forall g \in \widetilde{Aut}(R)). \tag{2.24}$$

We define $\mathbb{C}^*$-quotient spaces

$$\mathbb{P}(D^2) := D^2/\mathbb{C}^*, \tag{2.25}$$

$$\mathbb{P}(\mathbb{H}_{half}) := \mathbb{H}_{half}/\mathbb{C}^*. \tag{2.26}$$

These spaces are complex manifolds and they have $\widetilde{Aut}(R)$-actions. The $\mathbb{C}^*$-bundles $p : D^2 \to \mathbb{P}(D^2)$, $p : \mathbb{H}_{half} \to \mathbb{P}(\mathbb{H}_{half})$ are $\widetilde{Aut}(R)$-equivariant.

We have a natural diagram:

$$\begin{tikzcd} D^2 \arrow[r] \arrow[d, "p"'] & \mathbb{H}_{half} \arrow[d, "p"] \\ \mathbb{P}(D^2) \arrow[r] & \mathbb{P}(\mathbb{H}_{half}) \end{tikzcd} \quad . \tag{2.27}$$

This diagram is a $\mathbb{C}^*$-equivariant morphism from $D^2 \to \mathbb{P}(D^2)$ to $\mathbb{H}_{half} \to \mathbb{P}(\mathbb{H}_{half})$. Thus this diagram is cartesian. Also this diagram is $\widetilde{Aut}(R)$-equivariant.

2.5 A tensor on the domain D^2

2.5.1 Definition of a holomorphic metric on D

First we define a holomorphic metric I_D on D. Since D is an open set of an affine space, there exists uniquely the holomorphic metric I_D on D:

$$I_D \in \Gamma(D, \Omega^1_D \otimes_{\mathcal{O}_D} \Omega^1_D) \tag{2.28}$$

characterized by the property that on each $p \in D$, it gives

$$I_{D,p}(\delta, \delta') := I_{(F^2_{\mathbb{C}})^*}(f_p(\delta), f_p(\delta')) \quad (\forall \delta, \delta' \in T_pD) \tag{2.29}$$

where $f_p : T_pD \xrightarrow{\sim} (F^2_{\mathbb{C}})^*$ is a canonical isomorphism.

By this definition, we have

$$\varphi(g)^* I_D = I_D \quad (\forall g \in \widetilde{Aut}(R)), \tag{2.30}$$

$$\psi(\alpha)^* I_D = \alpha^2 I_D \quad (\forall \alpha \in \mathbb{C}^*). \tag{2.31}$$

2.5.2 Definition of a tensor on D^2

We define a tensor on D^2. We define $j_1^* I_D \in \Gamma(D^2, \Omega^1_{D^2} \otimes_{\mathcal{O}_{D^2}} \Omega^1_{D^2})$ as a pull-back of $I_D \in \Gamma(D, \Omega^1_D \otimes_{\mathcal{O}_D} \Omega^1_D)$ by the inclusion $j_1 : D^2 \hookrightarrow D$.

Proposition 2.7. *There exists uniquely an element* $I_{D^2} \in \Gamma(D^2, p^*\Omega^1_{\mathbb{P}(D^2)} \otimes_{\mathcal{O}_{D^2}} p^*\Omega^1_{\mathbb{P}(D^2)})$ *such that its image by the morphism*

$$\Gamma(D^2, p^*\Omega^1_{\mathbb{P}(D^2)} \otimes_{\mathcal{O}_{D^2}} p^*\Omega^1_{\mathbb{P}(D^2)}) \to \Gamma(D^2, \Omega^1_{D^2} \otimes_{\mathcal{O}_{D^2}} \Omega^1_{D^2}) \tag{2.32}$$

is $j_1^* I_D$. *The tensor* I_{D^2} *gives a non-degenerate* $\mathcal{O}_{D^2}$*-symmetric bilinear form on* $p^*\Theta_{\mathbb{P}(D^2)}$.

Proof. For $x \in D^2$, the radical of $\mathbb{C}$-blinear form $(j_1^* I_D)_x : T_x D^2 \times T_x D^2 \to \mathbb{C}$ is just a kernel of $(p_*)_x : T_x D^2 \to T_{p(x)}\mathbb{P}(D^2)$. Thus $(j_1^* I_D)_x$ gives a non-degenerate $\mathbb{C}$-bilinear form on $T_x D^2 / \ker(p_*)_x$. Thereby we have a result. ☐

By this proposition, we define its dual

$$I^*_{D^2} \in \Gamma(D^2, p^*\Theta_{\mathbb{P}(D^2)} \otimes_{\mathcal{O}_{D^2}} p^*\Theta_{\mathbb{P}(D^2)}). \tag{2.33}$$

By a uniqueness of I_{D^2} and equations (2.30), (2.31), we have

$$\varphi(g)^* I^*_{D^2} = I^*_{D^2} \quad (\forall g \in \widetilde{Aut}(R)), \tag{2.34}$$

$$\psi(\alpha)^* I^*_{D^2} = \alpha^{-2} I^*_{D^2} \quad (\forall \alpha \in \mathbb{C}^*). \tag{2.35}$$

2.6 Definition of the analytic Weyl group quotient

2.6.1 Open subset of the domain

For the open subsets of D^2 and $\mathbb{P}(D^2)$, we define W^2-quotient space and a tensor on it.

We put

$$\overset{\circ}{D}{}^2 := \{ x \in D^2 \mid \langle \alpha, x \rangle \neq 0 \ (\forall \alpha \in R) \}, \tag{2.36}$$

$$\mathbb{P}(\overset{\circ}{D}{}^2) := \overset{\circ}{D}{}^2 / \mathbb{C}^*, \tag{2.37}$$

whose complement is a reflection hyperplanes. Then $p : \overset{\circ}{D}{}^2 \to \mathbb{P}(\overset{\circ}{D}{}^2)$ is a $\mathbb{C}^*$-bundle and the diagram

$$\begin{tikzcd}
\overset{\circ}{D}{}^2 \arrow[r] \arrow[d, "p"'] & \mathbb{H}_{half} \arrow[d, "p"] \\
\mathbb{P}(\overset{\circ}{D}{}^2) \arrow[r] & \mathbb{P}(\mathbb{H}_{half})
\end{tikzcd} \tag{2.38}$$

is $\widetilde{Aut}(R)$-equivariant and cartesian. Since $j_2 : \overset{\circ}{D}{}^2 \to D^2$ is an open immersion, $j_2^* I^*_{D^2}$ is an element of $\Gamma(\overset{\circ}{D}{}^2, p^*\Theta_{\mathbb{P}(\overset{\circ}{D}{}^2)} \otimes_{\mathcal{O}_{\overset{\circ}{D}{}^2}} p^*\Theta_{\mathbb{P}(\overset{\circ}{D}{}^2)})$. We put $I^*_{\overset{\circ}{D}{}^2} := j_2^* I^*_{D^2}$.

Proposition 2.8. *The group* W^2 *acts on* $\overset{\circ}{D}{}^2$ *and* $\mathbb{P}(\overset{\circ}{D}{}^2)$ *properly discontinuous and fixed point free.*

Proof. Since $p : \overset{\circ}{D}{}^2 \to \mathbb{P}(\overset{\circ}{D}{}^2)$ is W^2-equivariant, we should only prove this Proposition for $\mathbb{P}(\overset{\circ}{D}{}^2)$, which will be shown in Proposition 5.5. ☐

2.6.2 Definition of an analytic Weyl group quotient

Since W^2-action and $\mathbb{C}^*$-action are commutative, we have a $\mathbb{C}^*$-bundle $p : \overset{\circ}{D}{}^2 /W^2 \to \mathbb{P}(\overset{\circ}{D}{}^2)/W^2$. In the following diagram:

$$\begin{tikzcd}
\overset{\circ}{D}{}^2 \arrow[r, "j_3"'] \arrow[d, "p"'] & \overset{\circ}{D}{}^2 /W^2 \arrow[r] \arrow[d, "p"'] & \mathbb{H}_{half} \arrow[d, "p"] \\
\mathbb{P}(\overset{\circ}{D}{}^2) \arrow[r, "j_3"'] & \mathbb{P}(\overset{\circ}{D}{}^2)/W^2 \arrow[r] & \mathbb{P}(\mathbb{H}_{half})
\end{tikzcd} , \tag{2.39}$$

all morphisms are $\widetilde{Aut}(R)$-equivariant and both squares are cartesian.

2.6.3 A tensor on an analytic Weyl group quotient space

We define a tensor on $\overset{\circ}{D}{}^2 /W^2$. Since $I^*_{\overset{\circ}{D^2}}$ is W^2-invariant by (2.34) and $j_3 : \overset{\circ}{D}{}^2 \to \overset{\circ}{D}{}^2 /W^2$ is étale, there exists uniquely a tensor $I^*_{\overset{\circ}{D^2}/W^2} \in \Gamma(\overset{\circ}{D}{}^2 /W^2, p^*\Theta_{\mathbb{P}(\overset{\circ}{D^2})/W^2} \otimes_{\mathcal{O}_{\overset{\circ}{D^2}/W^2}} p^*\Theta_{\mathbb{P}(\overset{\circ}{D^2})/W^2})$ satisfying $j_3^* I^*_{\overset{\circ}{D^2}/W^2} = I^*_{\overset{\circ}{D^2}}$.

By a uniqueness of $I^*_{\overset{\circ}{D^2}/W^2}$ and equations (2.34), (2.35), we have

$$\varphi(g)^* I^*_{\overset{\circ}{D^2}/W^2} = I^*_{\overset{\circ}{D^2}/W^2} \qquad (\forall g \in \widetilde{Aut}(R)), \tag{2.40}$$

$$\psi(\alpha)^* I^*_{\overset{\circ}{D^2}/W^2} = \alpha^{-2} I^*_{\overset{\circ}{D^2}/W^2} \qquad (\forall \alpha \in \mathbb{C}^*). \tag{2.41}$$

3 Algebraic Weyl group quotient spaces

3.1 Definition of Weyl group invariant rings

3.1.1 Definition of Weyl group invariants

In this subsection, we define the Weyl group invariant ring for the oriented elliptic root system with $\Lambda_{\mathbb{Z}} \neq \emptyset$.

By the diagram (2.27), we have the morphism:

$$\pi : D^2 \to \mathbb{P}(\mathbb{H}_{half}). \tag{3.1}$$

We define an $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-modules. For $k, m \in \mathbb{Z}$, we put

$$\begin{aligned}\boldsymbol{\mathcal{S}}^W_{D^2,k,m}(U) := \{ f \in \pi_*\mathcal{O}_{D^2}(U) | & \varphi(g)^* f = f \; (\forall g \in W^2), \\ & \varphi(\rho(t))^* f = e^{-2\pi i m t} f \; (\forall t \in \mathbb{C}), \\ & \psi(\alpha)^* f = \alpha^{-k} f \; (\forall \alpha \in \mathbb{C}^*) \, \},\end{aligned} \tag{3.2}$$

for an open set $U \subset \mathbb{P}(\mathbb{H}_{half})$. Here $\rho : \mathbb{C} \to K_{\mathbb{C}}$ is defined in Proposition 2.5.

3.1.2 Definition of Weyl group invariant rings

We define the $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-graded algebras by

$$\boldsymbol{\mathcal{S}}^W_{D^2,*,*} := \bigoplus_{k,m\in\mathbb{Z}} \boldsymbol{\mathcal{S}}^W_{D^2,k,m}, \;\; \boldsymbol{\mathcal{S}}^W_{D^2,*,0} := \bigoplus_{k\in\mathbb{Z}} \boldsymbol{\mathcal{S}}^W_{D^2,k,0}, \;\; \boldsymbol{\mathcal{S}}^W_{D^2,0,*} := \bigoplus_{m\in\mathbb{Z}} \boldsymbol{\mathcal{S}}^W_{D^2,0,m}. \tag{3.3}$$

3.1.3 Automorphism group action on the Weyl group invariant rings

In this subsection, we define an $\widetilde{Aut}(R)$-action and a $\mathbb{C}^*$-action on the ringed spaces $(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,*,*})$ etc.

Since the group $\widetilde{Aut}(R)$ acts on D^2, we have

$$\mathcal{O}_{D^2} \to \varphi(g)_*\mathcal{O}_{D^2}, \quad f \mapsto f \circ \varphi(g) \tag{3.4}$$

for a local section f of $\mathcal{O}_{D^2}$. Taking a direct image by π, we have

$$\pi_*\mathcal{O}_{D^2} \to \pi_*\varphi(g)_*\mathcal{O}_{D^2} \simeq \varphi(g)_*\pi_*\mathcal{O}_{D^2}. \tag{3.5}$$

A pair $\varphi(g) : \mathbb{P}(\mathbb{H}_{half}) \to \mathbb{P}(\mathbb{H}_{half})$ and the morphism (3.5) gives a morphism of a ringed space:

$$\varphi(g) : (\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{D^2}) \to (\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{D^2}) \; (g \in \widetilde{Aut}(R)). \tag{3.6}$$

These give an $\widetilde{Aut}(R)$-action of a ringed space $(\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{D^2})$. Since we could restrict the morphism (3.5) to the subsheaves $\boldsymbol{\mathcal{S}}^W_{D^2,*,*}$, $\boldsymbol{\mathcal{S}}^W_{D^2,*,0}$, $\boldsymbol{\mathcal{S}}^W_{D^2,0,*}$, and $\boldsymbol{\mathcal{S}}^W_{D^2,0,0}$, we obtain $\widetilde{Aut}(R)$-actions on the ringed spaces

$$\begin{aligned}&(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,*,*}), \; (\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,*,0}), \\ &(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,0,*}), \; (\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,0,0})\end{aligned} \tag{3.7}$$

and we denote the action also by φ.

In a same manner we could also define $\mathbb{C}^*$-actions on these ringed spaces and we denote the action also by ψ.

By a parallel construction, we could also define $\widetilde{Aut}(R)$-actions and $\mathbb{C}^*$-actions on the ringed spaces

$$\begin{aligned}&(\mathbb{P}(\mathbb{H}_{half}), \mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}), \quad (\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\mathbb{H}_{half}}), \\ &(\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\mathbb{P}(\overset{\circ}{D}^2)/W^2}), \quad (\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\overset{\circ}{D}^2/W^2}).\end{aligned} \tag{3.8}$$

3.2 Structure of Weyl group invariant rings

In this subsection, we study the structure of the invariant ring.

3.2.1 A Chevalley's type theorem

Theorem 3.1. ([1], [2], [4], [7], [8], [14])
The $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-graded algebra $\boldsymbol{\mathcal{S}}^W_{D^2,0,*}$ is an $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-free algebra, i.e.

$$\boldsymbol{\mathcal{S}}^W_{D^2,0,*} = \mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}[s^1, \cdots, s^{n-1}] \tag{3.9}$$

for $s^j \in \Gamma(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,0,*})$ with $c^1 \geq c^2 \geq \cdots \geq c^{n-1} > 0$ and $n := l+2$. We remark that j of s^j is a suffix.

As we see in §5.6, we reduce its proof to the works [1], [2], [4], [7], [8], [14].

Corollary 3.2. 1. *We have a natural inclusion morphism $(\mathrm{rad} I_F)_{\mathbb{C}} \to \Gamma(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,-1,0})$.*

2. *Take $a \in \mathrm{rad} I_F \setminus \{0\}$. By 1, we regard $a \in \Gamma(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,-1,0})$. We also have $a^{-1} \in \Gamma(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,1,0})$. Then we have isomoprhisms:*

$$\boldsymbol{\mathcal{S}}^W_{D^2,*,*} \simeq \mathbb{C}[a, a^{-1}] \otimes_{\mathbb{C}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*}, \tag{3.10}$$

$$\boldsymbol{\mathcal{S}}^W_{D^2,*,0} \simeq \mathbb{C}[a, a^{-1}] \otimes_{\mathbb{C}} \boldsymbol{\mathcal{S}}^W_{D^2,0,0}. \tag{3.11}$$

3. *We have a canonical isomoprhism:*

$$\boldsymbol{\mathcal{S}}^W_{D^2,*,*} \simeq \boldsymbol{\mathcal{S}}^W_{D^2,*,0} \otimes_{\boldsymbol{\mathcal{S}}^W_{D^2,0,0}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*}. \tag{3.12}$$

4. *The $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-algebras $\boldsymbol{\mathcal{S}}^W_{D^2,*,*}$, $\boldsymbol{\mathcal{S}}^W_{D^2,*,0}$, $\boldsymbol{\mathcal{S}}^W_{D^2,0,*}$ are of finite presentation over $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$.*

Proof. For 1, we first see that an element of $(\mathrm{rad} I_F)_{\mathbb{C}}$ is regarded as a holomorphic function on $\mathbb{H}_{half}$. By the morphism $D^2 \to \mathbb{H}_{half}$, we have $(\mathrm{rad} I_F)_{\mathbb{C}} \to \Gamma(\mathbb{P}(\mathbb{H}_{half}), \pi_* \mathcal{O}_{D^2})$. We could easily check that it gives 1. Since $a \in \Gamma(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,-1,0})$ is a nowhere vanishing holomorphic function, we have $a^{-1} \in \Gamma(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,-1,0})$. By $a^k \boldsymbol{\mathcal{S}}^W_{D^2,k,m} \simeq \boldsymbol{\mathcal{S}}^W_{D^2,0,m}$, we have 2. For 3, we show that a natural morphism $\boldsymbol{\mathcal{S}}^W_{D^2,*,0} \otimes_{\boldsymbol{\mathcal{S}}^W_{D^2,0,0}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*} \to \boldsymbol{\mathcal{S}}^W_{D^2,*,*}$ is an isomorphism. By 2, we have $\boldsymbol{\mathcal{S}}^W_{D^2,*,0} \otimes_{\boldsymbol{\mathcal{S}}^W_{D^2,0,0}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*} \simeq (\mathbb{C}[a, a^{-1}] \otimes_{\mathbb{C}} \boldsymbol{\mathcal{S}}^W_{D^2,0,0}) \otimes_{\boldsymbol{\mathcal{S}}^W_{D^2,0,0}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*} \simeq \mathbb{C}[a, a^{-1}] \otimes_{\mathbb{C}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*} \simeq \boldsymbol{\mathcal{S}}^W_{D^2,*,*}$. The 4th assertion is a direct consequence of $\boldsymbol{\mathcal{S}}^W_{D^2,0,0} \simeq \mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$, 2 and Theorem 3.1. □

3.3 Relation between Weyl group invariant rings

We consider the spaces D^2, $\mathbb{H}_{half}$, $\overset{\circ}{D}{}^2$, $\overset{\circ}{D}{}^2 / W^2$, $\mathbb{P}(\overset{\circ}{D}{}^2)/W^2$ as $\mathbb{P}(\mathbb{H}_{half})$-objects and we denote the structure morphisms by π.

Proposition 3.3. *1. By the morphisms $D^2 \to \mathbb{H}_{half} \to \mathbb{P}(\mathbb{H}_{half})$, we regard the $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-modules $\pi_*\mathcal{O}_{\mathbb{H}_{half}}$ and $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$ as submodules of $\pi_*\mathcal{O}_{D^2}$. Then we have*

  (a) $\boldsymbol{\mathcal{S}}^W_{D^2,0,0} = \mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$,

  (b) $\boldsymbol{\mathcal{S}}^W_{D^2,*,0} \subset \pi_*\mathcal{O}_{\mathbb{H}_{half}}$.

2. By the morphisms $\overset{\circ}{D}{}^2 \to \overset{\circ}{D}{}^2/W^2 \to \mathbb{P}(\overset{\circ}{D}{}^2)/W^2$, we regard the $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-modules $\pi_*\mathcal{O}_{\overset{\circ}{D}{}^2/W^2}$ and $\pi_*\mathcal{O}_{\mathbb{P}(\overset{\circ}{D}{}^2)/W^2}$ as submodules of $\pi_*\mathcal{O}_{\overset{\circ}{D}{}^2}$. We also regard $\boldsymbol{\mathcal{S}}^W_{D^2,0,*}$ and $\boldsymbol{\mathcal{S}}^W_{D^2,*,*}$ as a submodules of $\pi_*\mathcal{O}_{\overset{\circ}{D}{}^2}$. Then we have

  (a) $\boldsymbol{\mathcal{S}}^W_{D^2,0,*} \subset \pi_*\mathcal{O}_{\mathbb{P}(\overset{\circ}{D}{}^2)/W^2}$,

  (b) $\boldsymbol{\mathcal{S}}^W_{D^2,*,*} \subset \pi_*\mathcal{O}_{\overset{\circ}{D}{}^2/W^2}$.

3. By the above inclusion morphisms, we have morphism of ringed spaces:

$$(\mathbb{P}(\mathbb{H}_{half}), \mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}) = (\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,0,0}), \tag{3.13}$$

$$(\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\mathbb{H}_{half}}) \to (\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,*,0}), \tag{3.14}$$

$$(\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\mathbb{P}(\overset{\circ}{D}{}^2)/W^2}) \to (\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,0,*}), \tag{3.15}$$

$$(\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\overset{\circ}{D}{}^2/W^2}) \to (\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,*,*}). \tag{3.16}$$

These morphisms are all $\widetilde{Aut}(R)$-equivariant and $\mathbb{C}^*$-equivariant.

Proof. In Theorem 3.1, $c^i > 0$ for $1 \leq i \leq n-1$. Thus we have 1(a). By Corollary 3.2(2) and $a, a^{-1} \in \Gamma(\mathbb{P}(\mathbb{H}_{half}), \pi_*\mathcal{O}_{\mathbb{H}_{half}})$, we have 1(b). For 2, we could check the conditions directly. For 3, it is O.K. because our definitions of group actions are induced from D^2 and $\overset{\circ}{D}{}^2$. □

3.4 Definition of Specan

We remind the notion of Specan and see some properties.

Let $\mathcal{C}$ be a category of $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-algebra which is of finite presentation, For an object $\mathcal{A}$ of the category $\mathcal{C}$, the analytic space $\operatorname{Specan}\mathcal{A}$ could be defined by [6]. It is characterized by a natural isomorphism:

$$\operatorname{Hom}_{(An)/\mathbb{P}(\mathbb{H}_{half})}(X, \operatorname{Specan}\mathcal{A}) \simeq \operatorname{Hom}_{\mathcal{O}_X}(f^*\mathcal{A}, \mathcal{O}_X)$$

for an object $f : X \to \mathbb{P}(\mathbb{H}_{half})$ of the category $(An)/\mathbb{P}(\mathbb{H}_{half})$, which is a category of an analytic space with the structure morphism to $\mathbb{P}(\mathbb{H}_{half})$.

Since there exists a canonical isomorphism: $\operatorname{Hom}_{\mathcal{O}_X}(f^*\mathcal{A}, \mathcal{O}_X) \simeq \operatorname{Hom}_{\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}}(\mathcal{A}, f_*\mathcal{O}_X)$, we have

$$\operatorname{Hom}_{(An)/\mathbb{P}(\mathbb{H}_{half})}(X, \operatorname{Specan}\mathcal{A}) \simeq \operatorname{Hom}_{\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}}(\mathcal{A}, f_*\mathcal{O}_X). \tag{3.17}$$

We denote the structure morphism $\operatorname{Specan}\mathcal{A} \to \mathbb{P}(\mathbb{H}_{half})$ by π.

3.5 Algebraic Weyl group quotient space

3.5.1 Definition of Algebraic Weyl group quotient space

We define the Algebraic Weyl group quotient spaces by

$$D^2//W^2 := \mathrm{Specan}\,\boldsymbol{\mathcal{S}}^W_{D^2,*,*}, \tag{3.18}$$

$$\mathbb{P}(D^2)//W^2 := \mathrm{Specan}\,\boldsymbol{\mathcal{S}}^W_{D^2,0,*}. \tag{3.19}$$

The groups $\widetilde{Aut}(R)$ and $\mathbb{C}^*$ act on the ringed spaces $(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,*,*})$ and $(\mathbb{P}(\mathbb{H}_{half}), \boldsymbol{\mathcal{S}}^W_{D^2,0,*})$ as we see in §3.1.3. Then a natural morphism $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ has a structure of $\mathbb{C}^*$-bundle. Also the group $\widetilde{Aut}(R)$ acts as

$$\begin{tikzcd}
D^2//W^2 \arrow[r, "\varphi(g)"] \arrow[d, "p"'] & D^2//W^2 \arrow[d, "p"'] \\
\mathbb{P}(D^2)//W^2 \arrow[r, "\varphi(g)"] \arrow[d] & \mathbb{P}(D^2)//W^2, \arrow[d] \\
\mathbb{P}(\mathbb{H}_{half}) \arrow[r, "\varphi(g)"] & \mathbb{P}(\mathbb{H}_{half})
\end{tikzcd} \tag{3.20}$$

for $g \in \widetilde{Aut}(R)$. We remark that $\varphi(g) : D^2//W^2 \to D^2//W^2$ and $\varphi(g) : \mathbb{P}(D^2)//W^2 \to \mathbb{P}(D^2)//W^2$ are not $\mathbb{P}(\mathbb{H}_{half})$-morphisms.

3.5.2 Relation to the analytic Weyl group quotient space

Proposition 3.4. *1. We have the canonical isomorphism*

$$\mathbb{H}_{half} \simeq \mathrm{Specan}\,\boldsymbol{\mathcal{S}}^W_{D^2,*,0}. \tag{3.21}$$

2. *We have a following natural diagram:*

$$\begin{tikzcd}
\overset{\circ}{D^2}/W^2 \arrow[r, "j_4"] \arrow[d, "p"'] & D^2//W^2 \arrow[r] \arrow[d, "p"'] & \mathbb{H}_{half} \arrow[d, "p"'] \\
\mathbb{P}(\overset{\circ}{D^2})/W^2 \arrow[r, "j_4"] & \mathbb{P}(D^2)//W^2 \arrow[r] & \mathbb{P}(\mathbb{H}_{half})
\end{tikzcd} . \tag{3.22}$$

   All morphism are $\widetilde{Aut}(R)$-equivariant and both squares are cartesian.

3. *The composite morphisms of row arrows coincide with morphisms in diagram (2.39).*

4. *In the diagram, the morphisms $j_4 : \overset{\circ}{D^2}//W^2 \to D^2//W^2$ and $j_4 : \mathbb{P}(\overset{\circ}{D^2})/W^2 \to \mathbb{P}(D^2)//W^2$ are open immersions and their images are open dense.*

5. *An element* $(\mathrm{rad} I_F)_{\mathbb{C}}$ *gives a* $\mathbb{C}^*$*-equivariant holomorphic function on* $\mathbb{H}_{half}$*. Thus it also gives a* $\mathbb{C}^*$*-equivariant holomorphic function on* $D^2//W^2$*.*

Proof. We show 1. By Proposition 3.3(1)(b), we have $\boldsymbol{\mathcal{S}}^W_{D^2,*,0} \subset \pi_*\mathcal{O}_{\mathbb{H}_{half}}$. Then we have a morphism $\mathbb{H}_{half} \to \mathrm{Specan}\,\boldsymbol{\mathcal{S}}^W_{D^2,*,0}$. Since it is a $\mathbb{C}^*$-equivariant morphism of the $\mathbb{C}^*$-bundles $\mathbb{H}_{half} \to \mathbb{P}(\mathbb{H}_{half})$ and $\mathrm{Specan}\,\boldsymbol{\mathcal{S}}^W_{D^2,*,0} \to \mathbb{P}(\mathbb{H}_{half})$, we see that it is an isomorphism. We show 2. By $\boldsymbol{\mathcal{S}}^W_{D^2,*,*} \simeq \boldsymbol{\mathcal{S}}^W_{D^2,*,0} \otimes_{\boldsymbol{\mathcal{S}}^W_{D^2,0,0}} \boldsymbol{\mathcal{S}}^W_{D^2,0,*}$ in Corollary 3.2(2) and 1, we see that the right hand side of the diagram is cartesian. We show 3. Since the morphism $X \to \mathbb{H}_{half}$ is determined by the induced mapping $(\mathrm{rad} I_F)_{\mathbb{C}} \to \Gamma(X, \mathcal{O}_X)$, we could check that the morphisms $\overset{\circ}{D}{}^2\ /W^2 \to \mathbb{H}_{half}$ in this diagram and (2.39) are same. Since the below row arrows of (3.22) are a $\mathbb{C}^*$-quotient of the above row arrows of (3.22), we have a result. We show 4. The morphisms $j_4 : \overset{\circ}{D}{}^2\ //W^2 \to D^2//W^2$ and $j_4 : \mathbb{P}(\overset{\circ}{D}{}^2)/W^2 \to \mathbb{P}(D^2)//W^2$ could be defined by $\boldsymbol{\mathcal{S}}^W_{D^2,0,*} \subset \pi_*\mathcal{O}_{\mathbb{P}(\overset{\circ}{D}{}^2)/W^2}$ and $\boldsymbol{\mathcal{S}}^W_{D^2,*,*} \subset \pi_*\mathcal{O}_{\overset{\circ}{D}{}^2/W^2}$ in Proposition 3.3(2)(b). By Proposition 5.6, $j_4 : \mathbb{P}(\overset{\circ}{D}{}^2)/W^2 \to \mathbb{P}(D^2)//W^2$ is an open immersion and its image is open dense. Thus $j_4 : \overset{\circ}{D}{}^2\ //W^2 \to D^2//W^2$ is also an open immersion and its image is open dense. The 5th assertion is a direct consequence of 2. □

3.6 A tensor on the algebraic quotient space

We assert an existence of a tensor on an algebraic quotient space $D^2//W^2$.

Proposition 3.5. *There exists uniquely an element*

$$I^*_{D^2//W^2} \in \Gamma(D^2//W^2, p^*\Theta_{\mathbb{P}(D^2)//W^2} \otimes_{\mathcal{O}_{D^2//W^2}} p^*\Theta_{\mathbb{P}(D^2)//W^2}) \tag{3.23}$$

such that

$$j_4^* I^*_{D^2//W^2} = I^*_{\overset{\circ}{D}{}^2/W^2}. \tag{3.24}$$

We give a proof in §5.8.

By a uniqueness of $I^*_{D^2//W^2}$ and equations (2.40), (2.41), we have

$$\varphi(g)^* I^*_{D^2//W^2} = I^*_{D^2//W^2} \quad (\forall g \in \widetilde{Aut}(R)), \tag{3.25}$$

$$\psi(\alpha)^* I^*_{D^2//W^2} = \alpha^{-2} I^*_{D^2//W^2} \quad (\forall \alpha \in \mathbb{C}^*). \tag{3.26}$$

4 Results

4.1 Frobenius

4.1.1 Definition of Frobenius manifold

We remind the definition of Frobenius manifold.

Definition 4.1. ([5, p.146, Def. 9.1]) A Frobenius manifold is a tuple $(M, \circ, e, E, J)$ where M is a complex manifold of dimension ≥ 1 with holomorphic metric J and multiplication $\circ$ on the tangent bundle, e is a global unit field and E is another global vector field, subject to the following conditions:

1. the metric is invariant under the multiplication, i.e., $J(X \circ Y, Z) = J(X, Y \circ Z)$ for local sections $X, Y, Z \in \Theta_M$,
2. (potentiality) the $(3,1)$-tensor $\nabla \circ$ is symmetric (here, ∇ is the Levi-Civita connection of the metric), i.e., $\nabla_X(Y \circ Z) - Y \circ \nabla_X(Z) - \nabla_Y(X \circ Z) + X \circ \nabla_Y(Z) - [X, Y] \circ Z = 0$, for local sections $X, Y, Z \in \Theta_M$,
3. the metric J is flat,
4. e is a unit field and it is flat, i.e. $\nabla e = 0$,
5. the Euler field E satisfies $Lie_E(\circ) = 1 \cdot \circ$ and $Lie_E(J) = D \cdot J$ for some $D \in \mathbb{C}$.

4.1.2 Definition of Intersection form

Definition 4.2. ([3, p.191]) For a Frobenius manifold $(M, \circ, e, E, J)$, we define an intersection form $I^* : \Omega_M^1 \times \Omega_M^1 \to \mathcal{O}_M$ by

$$I^*(\omega_1, \omega_2) = J(E, J^*(\omega_1) \circ J^*(\omega_2)) \tag{4.1}$$

where $J^* : \Omega_M^1 \to \Theta_M$ is the isomorphism induced by J.

4.2 Conformal Frobenius structure and the intersection form

4.2.1 Conformal Frobenius structure

We define a notion of conformal Frobenius manifold.

Definition 4.3. *Let $L \xrightarrow{p} M$ be a $\mathbb{C}^*$-bundle. Let $(\circ, e, E, J)$ be a tuple where*

$$\circ \in \Gamma(L,\, p^*\Omega_M^1 \otimes_{\mathcal{O}_L} p^*\Omega_M^1 \otimes_{\mathcal{O}_L} p^*\Theta_M), \tag{4.2}$$

$$e \in \Gamma(L,\, p^*\Theta_M), \tag{4.3}$$

$$E \in \Gamma(L,\, p^*\Theta_M), \tag{4.4}$$

$$J \in \Gamma(L,\, p^*\Omega_M^1 \otimes_{\mathcal{O}_L} p^*\Omega_M^1). \tag{4.5}$$

We call a tuple $(\circ, e, E, J)$ a conformal Frobenius structure of $p : L \to M$ if it satisfies the following conditions:

1. *For $\mathbb{C}^*$-action, we have*

$$\psi(\alpha)^* \circ = \circ,\ \psi(\alpha)^* e = e,\ \psi(\alpha)^* E = E,\ \psi(\alpha)^* J = \alpha^2 J \tag{4.6}$$

   where $\psi(\alpha) : L \to L$ is a $\mathbb{C}^*$-action for $\alpha \in \mathbb{C}^*$.

2. *J is non-degenerate, i.e. J gives the isomorphism*

$$J : p^*\Theta_M \xrightarrow{\sim} p^*\Omega^1_M. \tag{4.7}$$

3. *There exists an open covering $M = \cup_{\lambda\in\Lambda} M_\lambda$ and sections $\iota_\lambda : M_\lambda \to p^{-1}(M_\lambda)$ such that a tuple $(M_\lambda, \iota^*_\lambda\circ, \iota^*_\lambda e, \iota^*_\lambda E, \iota^*_\lambda J)$ is a Frobenius manifold for each $\lambda \in \Lambda$ under the identifications*

$$\iota^*_\lambda p^*\Omega^1_M \simeq \Omega^1_{M_\lambda}, \quad \iota^*_\lambda p^*\Theta_M \simeq \Theta_{M_\lambda}. \tag{4.8}$$

4.2.2 A good section

We deine a notion of a good section.

Definition 4.4. *Let $p : L \to M$ be a $\mathbb{C}^*$-bundle and $(\circ, e, E, J)$ be a conformal Frobenius structure. Let $U \subset M$ be an open set and $\iota : U \to p^{-1}(U)$ be a section. Then we call (U, ι) a good section if $(U, \iota^*\circ, \iota^*e, \iota^*E, \iota^*J)$ is a Frobenius manifold.*

4.2.3 Intersection form

Definition 4.5. *We define the intersection form $I^* \in \Gamma(L, p^*\Theta_M \otimes_{\mathcal{O}_L} p^*\Theta_M)$ of a conformal Frobenius structure $(\circ, e, E, J)$ by*

$$I^*(\omega_1, \omega_2) = J(E, J^*(\omega_1) \circ J^*(\omega_2)), \tag{4.9}$$

where we denote J^* as an inverse of J defined in (4.7).

We remark that for a good section (U, ι), ι^*I^* gives an intersection form for a Frobenius manifold $(U, \iota^*\circ, \iota^*e, \iota^*E, \iota^*J)$.

4.3 Sufficient condition

We prepare a sufficient condition for the existence of a conformal Frobenius structure with a given intersection form.

We first fix some notations. For a trivial $\mathbb{C}^*$-bundle $p : L \to M$, the following data are equivalent:

1. $L \simeq M \times \mathbb{C}^*$.
2. $\iota : M \to L$ satisfying $p \circ \iota = id$.
3. $\mathbb{C}^*$-equivariant morphism $f : L \to \mathbb{C}^*$.

These are related as $M \times \mathbb{C}^* \simeq L,\ (x, \alpha) \mapsto \alpha \cdot \iota(x)$ and $L \simeq M \times \mathbb{C}^*,\ y \mapsto (p(x), f(y))$. We denote by $\iota(f) : M \to L$ a section which corresponds to a $\mathbb{C}^*$-equivariant morphism $f : L \to \mathbb{C}^*$.

We denote

$$V_M := \Gamma(M, (\Theta_M)^{\otimes l} \otimes_{\mathcal{O}_M} (\Omega^1_M)^{\otimes m}), \tag{4.10}$$

$$V_L := \Gamma(L, (p^*\Theta_M)^{\otimes l} \otimes_{\mathcal{O}_L} (p^*\Omega^1_M)^{\otimes m}), \tag{4.11}$$

$$V_L^k := \{\omega \in V_L \mid \psi(\alpha)^*\omega = \alpha^k \omega\}. \tag{4.12}$$

We put

$$\Psi : V_M \to V_L, \quad \omega \mapsto p^*\omega \tag{4.13}$$

be a natural morphism. The following lemma is easy.

Lemma 4.6. *Let $p : L \to M$ be a trivial $\mathbb{C}^*$-bundle.*

1. *Ψ induces an isomorphism $V_M \simeq V_L^0$.*
2. *Let $\iota(f)$ be a section which corresponds to a $\mathbb{C}^*$-equivariant morphism $f : L \to \mathbb{C}^*$. Then for any $k \in \mathbb{Z}$, the following isomorphisms*

$$V_M \simeq V_L^k, \quad \omega \mapsto f^k p^*\omega, \tag{4.14}$$

$$V_L^k \simeq V_M, \quad \omega \mapsto \iota(f)^*\omega \tag{4.15}$$

are inverse to each other.

3. *Let $g : L \to \mathbb{C}^*$ be a $\mathbb{C}^*$-equivariant morphism. Then for $\omega \in V_L^k$, we have*

$$\iota(g)^*\omega = (f/g)^k \iota(f)^*\omega, \tag{4.16}$$

where we regard f/g as a holomorphic function on M.

By a correspondence given in Lemma 4.6, we have a following sufficient condition for a existence of a conformal Frobenius manifold with an intersection form.

Proposition 4.7. *Let $p : L \to M$ be a trivial $\mathbb{C}^*$-bundle. Let $\iota(f)$ be a section which corresponds to a $\mathbb{C}^*$-equivariant morphism $f : L \to \mathbb{C}^*$. Let I^* be an element of $\Gamma(L, p^*\Theta_M \otimes_{\mathcal{O}_L} p^*\Theta_M)$ with $\psi^*(\alpha)I^* = \alpha^{-2}I^*$ for $\alpha \in \mathbb{C}^*$. We assume that $(M, \circ, e, E, J)$ is a Frobenius manifold with an intersection form ι^*I^*. Then $(p^*\circ, p^*e, p^*E, f^2p^*J)$ gives a conformal Frobenius structure with an intersection form I^*.*

4.4 Existence of a conformal Frobenius structure

We assert the existence of a conformal Frobenius structure of $p : D^2//W^2 \to \mathbb{P}(D^2)/W^2$ with suitable conditions.

We first define a Euler operator on $\mathbb{P}(D^2)//W^2$. For a local section f of $\mathcal{O}_{\mathbb{P}(D^2)//W^2}$, we put

$$E_{\mathbb{P}(D^2)//W^2} f := \frac{-1}{c^1} \frac{1}{2\pi\sqrt{-1}} \lim_{t \to 0} \frac{\varphi(\rho(t))^*f - f}{t}. \tag{4.17}$$

Then $E_{\mathbb{P}(D^2)//W^2}$ gives a holomorphic vector field on $\mathbb{P}(D^2)$. For the oriented elliptic root system with $\Lambda_{\mathbb{Z}} \neq \emptyset$, the condition $c^1 > c^2$ is called "codimension 1" in [11], where c^1 and c^2 are defined in Theorem 3.1.

Theorem 4.8. *If the oriented elliptic root system with $\Lambda_{\mathbb{Z}} \neq \emptyset$ satisfies the condition of codimension 1, then there exists a conformal Frobenius structure $(\circ, e, E, J)$ of $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ satisfying the following properties:*

1. *Its intersection form coincides with $I^*_{D^2//W^2}$.*
2. *Its Euler field coincides with $E_{\mathbb{P}(D^2)//W^2}$ under the identification of Lemma 4.6(1).*

For a proof, we give it in §5.9.

4.5 Uniqueness of a conformal Frobenius structure

We assert that a conformal Frobenius structure of $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ is unique up to $\mathbb{C}^*$ multiplication under some conditions.

Theorem 4.9. *Let $(\circ, e, E, J)$, $(\circ', e', E, J')$ be conformal Frobenius structures of $p : D^2//W^2 \to \mathbb{P}(D^2)/W^2$ which satisfies conditions of 1 and 2 of Theorem 4.8. We also assume that they have global good sections. Then there exists $c \in \mathbb{C}^*$ such that*

$$(\circ', e', E, J') = (c^{-1}\circ, ce, E, c^{-1}J). \tag{4.18}$$

For a proof, we give it in §5.11.

4.6 Good sections

We classify good sections of a conformal Frobenius structure of Theorem 4.8.

Proposition 4.10. *We fix a conformal Frobenius structure of $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$.*

1. *For any $f \in (\mathrm{rad} I_F)_{\mathbb{C}} \setminus \{0\}$, it defines $f : D^2//W^2 \setminus \{f = 0\} \to \mathbb{C}^*$ and the corresponding section $\iota(f) : \mathbb{P}(D^2)//W^2 \setminus p(\{f = 0\}) \to D^2//W^2 \setminus \{f = 0\}$. Then $\iota(f)$ is a good section.*
2. *Let (U, ι) be a good section of this conformal Frobenius structure. Then the exists uniquely an element $f \in (\mathrm{rad} I_F)_{\mathbb{C}} \setminus \{0\}$ such that $\iota = \iota(f)|_U$.*

For a proof, we give it in §5.12.

4.7 The automorphism group action

We discuss a transformation of a conformal Frobenius structure by the action of $\widetilde{Aut}(R)$ on $D^2//W^2$.

Proposition 4.11. *We fix a conformal Frobenius structure* $(\circ, e, E, J)$ *of* $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$*. Then there exists the group homomorphism*

$$\chi : \widetilde{Aut}(R) \to \mathbb{C}^* \tag{4.19}$$

such that

$$(\varphi(g)^*\circ, \varphi(g)^*e, \varphi(g)^*E, \varphi(g)^*J) = (\chi(g)^{-1}\circ, \chi(g)e, E, \chi(g)^{-1}J) \tag{4.20}$$

for any $g \in \widetilde{Aut}(R)$*.*

Proof. We fix $g \in \widetilde{Aut}(R)$. We see that $(\varphi(g)^*\circ, \varphi(g)^*e, \varphi(g)^*E, \varphi(g)^*J)$ is a conformal Frobenius structure of $p : D^2//W^2 \to \mathbb{P}(D^2)//W^2$ with intersection form $\varphi(g)^*I^*_{D^2//W^2}$. By (3.25), we have $\varphi(g)^*I^*_{D^2//W^2} = I^*_{D^2//W^2}$. Since $E = E_{\mathbb{P}(D^2)//W^2}$ and $E_{\mathbb{P}(D^2)//W^2}$ is a center of $\widetilde{Aut}(R)$, we have $\varphi(g)^*E = E$. Then by a uniqueness of Theorem 4.9, we see that

$$(\varphi(g)^*\circ, \varphi(g)^*e, \varphi(g)^*E, \varphi(g)^*J) = (c(g)^{-1}\circ, c(g)e, E, c(g)^{-1}J). \tag{4.21}$$

for some $c(g) \in \mathbb{C}^*$. Since $\varphi(g)^*e = c(g)e$ and $e \neq 0$, $c(g)$ is uniquely determined and $c(g)$ satisfies $c(gg') = c(g)c(g')$. We put $\chi(g) := c(g)$. Then we have a result. □

5 Proofs

5.1 Domains for a signed marking

Hereafter we fix an element $a \in \Lambda_{\mathbb{Z}}$. We take $b \in \mathrm{rad} I_F$ such that a, b is an oriented basis. We put

$$D_a := \{x \in D \mid \langle a, x\rangle = 1\}, \tag{5.1}$$

$$D^2_a := D_a \cap D^2, \tag{5.2}$$

$$\mathbb{H}_a := \{x \in \mathbb{H}_{half} \mid \langle a, x\rangle = 1\}, \tag{5.3}$$

$$F^a_{\mathbb{C}} := \{x \in F^2_{\mathbb{C}} \mid I_{F^2_{\mathbb{C}}}(a, x) = 0\}, \tag{5.4}$$

$$(F^a_{\mathbb{C}})^* := \mathrm{Hom}_{\mathbb{C}}(F^a_{\mathbb{C}}, \mathbb{C}), \tag{5.5}$$

$$D^1_a := \{x \in (F^a_{\mathbb{C}})^* \mid \langle a, x\rangle = 1,\ \mathrm{Im}\langle b, x\rangle > 0\}. \tag{5.6}$$

The spaces D_a, D^2_a and D^1_a are complex manifolds with natural W^2-action and $K_{\mathbb{C}}$-action. The space D^1_a does not depend on the choice of $b \in \mathrm{rad} I_F$.

Remark 5.1. *We remark that* $F^a_{\mathbb{C}}$ *gives a complexification of a hyperbolic extension defined in [11]. Thus* D^1_a *corresponds to a space* $\tilde{\mathbb{E}}$ *defined in [11].*

We have a natural diagram:

$$\begin{tikzcd}
D_a^1 & D_a \arrow[l, "f_2"'] \arrow[d, "i"'] & D_a^2 \arrow[l, "f_1"'] \arrow[r] \arrow[d, "i"'] & \mathbb{H}_a \arrow[d, "i"'] \\
& D & D^2 \arrow[l, "j_1"'] \arrow[r] \arrow[d, "p"'] & \mathbb{H}_{half} \arrow[d, "p"] \\
& & \mathbb{P}(D^2) \arrow[r] & \mathbb{P}(\mathbb{H}_{half})
\end{tikzcd} \quad . \tag{5.7}$$

All three squares are cartesian. This diagram is W^2-equivariant and $K_{\mathbb{C}}$-equivariant. We also remark that the morphisms

$$f_2 \circ f_1 : D_a^2 \to D_a^1, \quad p \circ i : D_a^2 \to \mathbb{P}(D^2) \tag{5.8}$$

are isomorphisms.

5.2 A tensor on the domain

We define holomorphic metrics on D_a^2 and D_a^1. We put

$$I_{D_a^2} := i^* j_1^* I_D \in \Gamma(D_a^2, \Omega^1_{D_a^2} \otimes_{\mathcal{O}_{D_a^2}} \Omega^1_{D_a^2}), \tag{5.9}$$

$$I_{D_a^1} := (f_2 \circ f_1)^* I_{D_a^2} \in \Gamma(D_a^1, \Omega^1_{D_a^1} \otimes_{\mathcal{O}_{D_a^1}} \Omega^1_{D_a^1}). \tag{5.10}$$

By a proof of Proposition 2.7, $I_{D_a^2}$ is non-degenerate. Since $f_2 \circ f_1$ is an isomorphism, $I_{D_a^1}$ is also non-degenerate. We define their duals:

$$I^*_{D_a^2} \in \Gamma(D_a^2, \Theta_{D_a^2} \otimes_{\mathcal{O}_{D_a^2}} \Theta_{D_a^2}), \tag{5.11}$$

$$I^*_{D_a^1} \in \Gamma(D_a^1, \Theta_{D_a^1} \otimes_{\mathcal{O}_{D_a^1}} \Theta_{D_a^1}). \tag{5.12}$$

We remark that $I^*_{D_a^1}$ is W^2-invariant. The following Proposition gives a characterization of the tensor $I^*_{D_a^1}$.

Proposition 5.2. *For any $x \in D_a^1$, we have a canonical isomorphism $T_x^* D_a^1 \simeq F_{\mathbb{C}}^a/\mathbb{C}a$ and we have a commutative diagram:*

$$\begin{tikzcd}
(I^*_{D_a^1})_x : & T_x^* D_a^1 & \times & T_x^* D_a^1 \arrow[r] & \mathbb{C} \\
I_{F_{\mathbb{C}}^a/\mathbb{C}a} : & F_{\mathbb{C}}^a/\mathbb{C}a \arrow[u] & \times & F_{\mathbb{C}}^a/\mathbb{C}a \arrow[u] \arrow[r] & \mathbb{C} \arrow[u, equal]
\end{tikzcd}, \tag{5.13}$$

where $I_{F_{\mathbb{C}}^a/\mathbb{C}a}$ is a $\mathbb{C}$-bilinear mapping induced from $I_{F_{\mathbb{C}}^2}$.

Proof. Take $y \in D_a^2$ such that $f_2 \circ f_1(y) = x$. Then we have

$$\begin{tikzcd}
T_x D_a^1 & T_{f_1(y)} D_a \arrow[l, "(f_2)_*"'] \arrow[d, "i_*"'] & T_y D_a^2 \arrow[l, "(f_1)_*"'] \\
& T_{i \circ f_1(y)} D &
\end{tikzcd} \tag{5.14}$$

and a canonical isomorphisms:

$$\begin{tikzcd}
T_x D_a^1 & T_{f_1(y)} D_a \arrow[l, "(f_2)_*"'] \arrow[r, "i_*"] & T_{i \circ f_1(y)} D \\
(F_{\mathbb{C}}^a / \mathbb{C}a)^* \arrow[u, "\wr"] & (F_{\mathbb{C}}^2 / \mathbb{C}a)^* \arrow[l] \arrow[r] \arrow[u, "\wr"] & (F_{\mathbb{C}}^2)^* \arrow[u, "\wr"]
\end{tikzcd} . \tag{5.15}$$

From this, we have

$$\ker[(f_2)_* : T_{f_1(y)} D_a \to T_x D_a^1] = \mathrm{rad}(i^* I_D)_{f_1(y)}. \tag{5.16}$$

Then $(i^* I_D)_{f_1(y)}$ induces a $\mathbb{C}$-bilinear form on $T_x D_a^1$. We see easily that it coincides with $I_{D_a^1}$ because $I_{D_a^1}$ could be written as $(f_2 \circ f_1)^* f_1^* i^* I_D$. $\square$

Remark 5.3. *By this Proposition, we see that a tensor* $I_{D_a^1}^*$ *coincides with a tensor* $I_{\tilde{\mathbb{E}}}^*$ *defined in [11] under the identification* $D_a^1 \simeq \tilde{\mathbb{E}}$.

Remark 5.4. *We also comment of a relation to the work [12]. In the diagram (5.7), the group* $\widetilde{Aut}(R)$ *acts on* $p : D^2 \to \mathbb{P}(D^2)$. *But by* $g \in \widetilde{Aut}(R)$, D_a^2 *goes to* $D_{g(a)}^2$. *So we do not have a natural* $\widetilde{Aut}(R)$ *action on* D_a^2 *nor* D_a^1. *By the identification of* D_a^2 *and* D_a^1 *with* $\mathbb{P}(D^2)$, *we have an* $\widetilde{Aut}(R)$*-action on* D_a^2 *and* D_a^1. *Then the tensors* $I_{D_a^2}^*$ *and* $I_{D_a^1}^*$ *admit a conformal transformation by this action. This explains the reason of the appearance of a conformal transformation in the work of [12].*

5.3 Open subset of the domain and its analytic Weyl group quotient space

We put

$$\mathring{D}_a^2 := \{ x \in D_a^2 \mid \langle \alpha, x \rangle \neq 0 \ (\forall \alpha \in R) \}, \tag{5.17}$$

$$\mathring{D}_a^1 := \{ x \in D_a^1 \mid \langle \alpha, x \rangle \neq 0 \ (\forall \alpha \in R) \}. \tag{5.18}$$

The group W^2 acts on $\mathring{D}_a^2$, $\mathring{D}_a^1$ and we have a diagram:

$$\begin{tikzcd}
\mathring{D}_a^1 \arrow[d] & \mathring{D}_a^2 \arrow[l] \arrow[r] \arrow[d] & \mathbb{P}(\mathring{D}^2) \arrow[d] \\
D_a^1 & D_a^2 \arrow[l, "f_2 \circ f_1"'] \arrow[r, "p \circ i"] & \mathbb{P}(D^2)
\end{tikzcd} \tag{5.19}$$

which is W^2-equivariant.

Proposition 5.5. *The group W^2 acts on $\overset{\circ}{D}{}^1_a$, $\overset{\circ}{D}{}^2_a$, $\mathbb{P}(\overset{\circ}{D}{}^2)$ properly discontinuous and fixed point free.*

Proof. We need only to show it for $\overset{\circ}{D}{}^1_a$. By Remark (5.1), we should prove for the domain $\tilde{\mathbb{E}}$ in [11]. It is proved in [11]. $\square$

5.4 A tensor on the analytic Weyl group quotient

Since $j_3 : \overset{\circ}{D}{}^2_a \to \overset{\circ}{D}{}^2_a / W^2$ and $j_3 : \overset{\circ}{D}{}^1_a \to \overset{\circ}{D}{}^1_a / W^2$ are étale, there exist uniquely

$$I^*_{\overset{\circ}{D}{}^2_a/W^2} \in \Gamma(\overset{\circ}{D}{}^2_a / W^2, \Theta_{\overset{\circ}{D}{}^2_a/W^2} \otimes_{\mathcal{O}_{\overset{\circ}{D}{}^2_a/W^2}} \Theta_{\overset{\circ}{D}{}^2_a/W^2}), \tag{5.20}$$

$$I^*_{\overset{\circ}{D}{}^1_a/W^2} \in \Gamma(\overset{\circ}{D}{}^1_a / W^2, \Theta_{\overset{\circ}{D}{}^1_a/W^2} \otimes_{\mathcal{O}_{\overset{\circ}{D}{}^1_a/W^2}} \Theta_{\overset{\circ}{D}{}^1_a/W^2}) \tag{5.21}$$

such that

$$I^*_{\overset{\circ}{D}{}^2_a} = j_3^* I_{\overset{\circ}{D}{}^2_a/W^2}, \quad I^*_{\overset{\circ}{D}{}^1_a} = j_3^* I_{\overset{\circ}{D}{}^1_a/W^2}. \tag{5.22}$$

5.5 Weyl group invariant ring

By the diagram (5.7), we regard $\overset{\circ}{D}{}^2_a$, $\overset{\circ}{D}{}^1_a$ and $\mathbb{H}_a$ as $\mathbb{P}(\mathbb{H}_{half})$-objects and we denote the structure morphisms by π. We put

$$\boldsymbol{\mathcal{S}}^W_{D^2_a,m}(U) := \{ f \in \pi_* \mathcal{O}_{D^2_a}(U) \mid \varphi(g)^* f = f \ (\forall g \in W^2), \tag{5.23}$$
$$\varphi(\rho(t))^* f = e^{-2\pi i m t} f \ (\forall t \in \mathbb{C}) \},$$

$$\boldsymbol{\mathcal{S}}^W_{D^1_a,m}(U) := \{ f \in \pi_* \mathcal{O}_{D^1_a}(U) \mid \varphi(g)^* f = f \ (\forall g \in W^2), \tag{5.24}$$
$$\varphi(\rho(t))^* f = e^{-2\pi i m t} f \ (\forall t \in \mathbb{C}) \},$$

$$\boldsymbol{\mathcal{S}}^W_{\mathbb{H}_a} := \pi_* \mathcal{O}_{\mathbb{H}_a} \tag{5.25}$$

for an open set $U \subset \mathbb{P}(\mathbb{H}_{half})$. We define the $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-graded algebras by

$$\boldsymbol{\mathcal{S}}^W_{D^2_a,*} := \oplus_{m \in \mathbb{Z}} \boldsymbol{\mathcal{S}}^W_{D^2_a,m}, \quad \boldsymbol{\mathcal{S}}^W_{D^1_a,*} := \oplus_{m \in \mathbb{Z}} \boldsymbol{\mathcal{S}}^W_{D^1_a,m}. \tag{5.26}$$

5.6 Reduction of a proof of Theorem 3.1 to the previous works

Since $\mathcal{O}_{\mathbb{P}(\mathbb{H}_{half})}$-graded algebras $\boldsymbol{\mathcal{S}}^W_{D^2_a,*}$, $\boldsymbol{\mathcal{S}}^W_{D^1_a,*}$ and $\boldsymbol{\mathcal{S}}^W_{D^2,0,*}$ are isomorphic, we need only to show Theorem 3.1 for $\boldsymbol{\mathcal{S}}^W_{D^1_a,*}$, which is already shown in [1], [2], [4], [7], [8], [14].

5.7 Algebraic Weyl group quotient spaces

We define the Weyl group quotient spaces by

$$D^2_a//W^2 := \operatorname{Specan} \boldsymbol{\mathcal{S}}^W_{D^2_a,*}, \tag{5.27}$$

$$D^1_a//W^2 := \operatorname{Specan} \boldsymbol{\mathcal{S}}^W_{D^1_a,*}. \tag{5.28}$$

These spaces are related to the spaces which we defined in the previous sections.

Proposition 5.6. *1. We have a canonical isomorphism:*

$$\mathbb{H}_a \simeq \operatorname{Specan} \boldsymbol{\mathcal{S}}^W_{\mathbb{H}_a}. \tag{5.29}$$

2. We have a following natural diagram:

$$\begin{tikzcd}
\overset{\circ}{D^1_a}/W^2 \arrow[r, "j_4"] & D^1_a//W^2 \arrow[r] & \mathbb{H}_a \\
\overset{\circ}{D^2_a}/W^2 \arrow[r, "j_4"] \arrow[u, "f_2\circ f_1"] \arrow[d, "i"'] & D^2_a//W^2 \arrow[r] \arrow[u, "f_2\circ f_1"] \arrow[d, "i"'] & \mathbb{H}_a \arrow[u, equal] \arrow[d, "i"'] \\
\overset{\circ}{D^2}/W^2 \arrow[r, "j_4"] \arrow[d, "p"'] & D^2//W^2 \arrow[r] \arrow[d, "p"'] & \mathbb{H}_{half} \arrow[d, "p"'] \\
\mathbb{P}(\overset{\circ}{D^2})/W^2 \arrow[r, "j_4"] & \mathbb{P}(D^2)//W^2 \arrow[r] & \mathbb{P}(\mathbb{H}_{half})
\end{tikzcd} . \tag{5.30}$$

These squares are all cartesian.

3. j_4 are all open immersion and the image is open dense.

4. By Proposition 3.4(5), $a \in \Lambda_{\mathbb{Z}}$ defines a $\mathbb{C}^*$-equivariant morphism $a : D^2//W^2 \to \mathbb{C}^*$. By a correspondence in §4.3 it defines a section $\iota(a) : \mathbb{P}(D^2)//W^2 \to D^2//W^2$. We have $\iota(a) = i \circ (p \circ i)^{-1}$.

Proof. A proof of 1,2 are parallel to Proposition 3.4. For a proof of 3, we should only prove it for $j_4 : \overset{\circ}{D^1_a}/W^2 \to D^1_a//W^2$. By the identification in Remark 5.1, we reduce it to the corresponding result, which is shown in [11]. See also (3.12) and below of [13]. The 4th assertion is a direct consequence of 2. □

5.8 Proof of Proposition 3.5

Proposition 5.7. *There exists uniquely*

$$I^*_{D^2_a//W^2} \in \Gamma(D^2_a//W^2, \Theta_{D^2_a//W^2} \otimes_{\mathcal{O}_{D^2_a//W^2}} \Theta_{D^2_a//W^2}), \tag{5.31}$$

$$I^*_{D^1_a//W^2} \in \Gamma(D^1_a//W^2, \Theta_{D^1_a//W^2} \otimes_{\mathcal{O}_{D^1_a//W^2}} \Theta_{D^1_a//W^2}) \tag{5.32}$$

such that

$$j_4^* I^*_{D_a^2//W^2} = I^*_{\overset{\circ}{D_a^2}/W^2}, \quad j_4^* I^*_{D_a^1//W^2} = I^*_{\overset{\circ}{D_a^1}/W^2}. \tag{5.33}$$

Proof. For the existence of $I^*_{D_a^1//W^2}$, we proved in [13] under the identification in Remark 5.1, 5.3. We put

$$I^*_{D_a^2//W^2} := (f_2 \circ f_1)^* I^*_{D_a^1//W^2}. \tag{5.34}$$

Then we have the result. □

We define a tensor $I^*_{D^2//W^2}$ by

$$I^*_{D^2//W^2} := a^{-2} p^*((p \circ i)^{-1})^* I^*_{D_a^2//W^2}. \tag{5.35}$$

Then we have

$$\begin{aligned} j_4^* I^*_{D^2//W^2} &= j_4^* a^{-2} p^*((p \circ i)^{-1})^* I^*_{D_a^2//W^2} \\ &= a^{-2} p^*((p \circ i)^{-1})^* j_4^* I^*_{D_a^2//W^2} \\ &= a^{-2} p^*((p \circ i)^{-1})^* I^*_{\overset{\circ}{D^2}{}_a/W^2} \\ &= a^{-2} p^*((p \circ i)^{-1})^* i^* I^*_{\overset{\circ}{D^2}/W^2} \\ &= a^{-2} p^*(i \circ (p \circ i)^{-1})^* I^*_{\overset{\circ}{D^2}/W^2} \\ &= a^{-2} p^* \iota(a)^* I^*_{\overset{\circ}{D^2}/W^2} \\ &= I^*_{\overset{\circ}{D^2}/W^2}, \end{aligned}$$

where we use the correspondence in Lemma 4.6 for the last equality.

Since $\overset{\circ}{D^2}\ //W^2$ is open dense in $D^2//W^2$, the uniqueness of $I^*_{D^2//W^2}$ is apparent. Thus we obtain a proof of the Proposition 3.5. From the definition of $I^*_{D^2//W^2}$, we have the following easy consequence.

Corollary 5.8. *By the identification* $i^* p^* \Theta_{\mathbb{P}(D^2)//W^2} \simeq \Theta_{D_a^2//W^2}$*, we have*

$$i^* I^*_{D^2//W^2} = I^*_{D_a^2//W^2}. \tag{5.36}$$

5.9 Proof of Theorem 4.8

. By the identification in Remark 5.1 and 5.3, we have a following result by [13].

Theorem 5.9. *The space* $D_a^1//W^2$ *has a structure of Frobenius manifold* $(D_a^1//W^2, \circ, e, E, J)$ *whose intersection form is* $I^*_{D_a^2//W^2}$ *and* $E = E_{D_a^1//W^2}$*, where we define* $E_{D_a^1//W^2}$ *by*

$$E_{D_a^1//W^2} f := \frac{-1}{c^1} \frac{1}{2\pi\sqrt{-1}} \lim_{t \to 0} \frac{\varphi(\rho(t))^* f - f}{t} \tag{5.37}$$

for a local section $f \in \mathcal{O}_{D_a^1//W^2}$*. We also assert that*

1. *$J(e) \in \mathbb{C}^* d(b/a)$,*
2. *$d(b/a)$ is flat for $I^*_{D^1_a//W^2}$*

where we regard b/a as a holomorphic function on $D^1_a//W^2$.

Here we identify the operator $E_{D^1_a//W^2}$ with the operator in [13] by the Remark 2.6.

By the isomorphims

$$D^1_a//W^2 \xleftarrow{f_2 \circ f_1} D^2_a//W^2 \xrightarrow{p \circ i} \mathbb{P}(D^2)//W^2 \tag{5.38}$$

in the diagram (5.30), we introduce the structure of Frobenius manifold on $\mathbb{P}(D^2)//W^2$. Its Euler field is $E_{\mathbb{P}(D^2)//W^2}$ because the diagram (5.30) is $K_{\mathbb{C}}$-equivariant. Its intersection form is

$$\begin{aligned}
&\{(p \circ i)^{-1}\}^*(f_2 \circ f_1)^* I^*_{D^1_a//W^2} \\
=&\{(p \circ i)^{-1}\}^* I^*_{D^2_a//W^2} \qquad (\because\ (5.34)) \\
=&\{(p \circ i)^{-1}\}^* i^* I^*_{D^2//W^2} \qquad (\because \text{Corollary 5.8}) \\
=&\iota(a)^* I^*_{D^2//W^2} \qquad (\because \text{Proposition 5.6(4)}).
\end{aligned}$$

Then by Proposition 4.7, we have a proof of Theorem 4.8.

5.10 Deformation of holomorphic flat metric

We give two Propositions. First is well-known and could be proved directly.

Proposition 5.10. *Let M be a complex manifold with dimension $n \geq 3$. Let g, g' be flat holomorphic metrics on the complex manifold M satisfying*

$$g' = \sigma^2 g \tag{5.39}$$

for a nowhere-vanishing holomorphic function σ. Then σ must be of the form

1. *σ^{-1} is constant,*
2. *$\sigma^{-1} = \sum_{ij} g_{ij}(x^i - c^i)(x^j - c^j)$ for a flat coordinate $x^1, \cdots, x^n$ with respect to g and c^i $(1 \leq i \leq n)$*
3. *$\sigma^{-1} = c - \sum_i b_i x^i$ for $c, b_i \in \mathbb{C}$ $(1 \leq i \leq n)$ satisfying $g^*(d\sigma^{-1}, d\sigma^{-1}) = 0$ for a dual metric of g.*

The following proposition is a Frobenius manifold version of the above proposition.

Proposition 5.11. *Let $(M, \circ, e, E, J)$ be a Frobenius manifold with* $\dim M \geq 3$, *$J(e,e) = 0$. Let $\sigma : M \to \mathbb{C}^*$ be a holomorphic function.*
(1) $(M, \circ, e, E, J_\sigma := \sigma^2 J)$ is a Frobenius manifold if and only if σ is homogeneous with respect to E and $d(\sigma^{-1}) = cJ(e)$ for some $c \in \mathbb{C}$.
(2) If $Lie_E J = D \cdot J$, then $Lie_E(\sigma^2 J) = (2 - D)\sigma^2 J$.

Proof. We give a proof "if" part of (1). We check the conditions of Frobenius manifold. For a property of multiplication invariance of metric :

$$J_\sigma(X \circ Y, Z) = J_\sigma(X, Y \circ Z), \tag{5.40}$$

it is O.K. by $J(X \circ Y, Z) = J(X, Y \circ Z)$.

For a property of potentiality, we should prove that

$$\nabla^\sigma_X(Y \circ Z) - \nabla^\sigma_Y(X \circ Z) + X \circ \nabla^\sigma_Y(Z) - Y \circ \nabla^\sigma_X(Z) - [X, Y] \circ Z = 0 \tag{5.41}$$

for the Levi-Civita connection ∇^σ for a metric J_σ.

Let ∇ be the the Levi-Civita connection for a metric J. Since the relation of ∇^σ and ∇ are known as

$$\nabla^\sigma_X(Y) - \nabla_X(Y) = \alpha(X)Y + (X \wedge U)(Y) \tag{5.42}$$

where

$$\alpha := \frac{d\sigma}{\sigma},\ U := J(\alpha),\ (X \wedge Y)(Z) := J(Y, Z)X - J(X, Z)Y. \tag{5.43}$$

The equation (5.41) holds if we replace ∇^σ with ∇. Then we have

$$\begin{aligned}
&\text{LHS of (5.41)}\\
=&\alpha(X)(Y \circ Z) + (X \wedge U)(Y \circ Z) - \alpha(Y)(X \circ Z) - (Y \wedge U)(X \circ Z)\\
&+ X \circ (\alpha(Y)Z) + X \circ [(Y \wedge U)(Z)] - Y \circ (\alpha(X)Z) - Y \circ [(X \wedge U)(Z)]\\
=&(X \wedge U)(Y \circ Z) - (Y \wedge U)(X \circ Z) + X \circ [(Y \wedge U)(Z)] - Y \circ [(X \wedge U)(Z)]\\
=&J(U, Y \circ Z)X - J(U, X \circ Z)Y - J(Y, Z)X \circ U + J(X, Z)Y \circ U.
\end{aligned}$$

Here we used the property that a product $\circ$ is function linear, commutative. We also used that a metric J is multiplication invariant.

Since $U = -c\,\sigma\,e$, we have $J(U, Y \circ Z)X = J(Y, Z)X \circ U$ and $J(U, X \circ Z)Y = J(X, Z)Y \circ U$. Thus we see that (5.41) holds.

For a property of flatness for J_σ, we remind the relation between the curvature tensor of J and J_σ. Let R (resp. R^σ) be a curvature tensor of J (resp. J_σ). Then

$$R^\sigma(X, Y) = R(X, Y) - (B(X) \wedge Y + X \wedge B(Y)), \tag{5.44}$$

where $B(X) := -\alpha(X)U + \nabla_X(U) + \frac{1}{2}\alpha(U)X$, with α, U are defined in (5.43). We have $R(X, Y) = 0$ since J is flat.

In order to show that $B(X) = 0$, we show that $\alpha(U) = 0$ and $-\alpha(X)U + \nabla_X(U) = 0$. The former part is as follows. $\alpha(U) = J(U, U) = c^2\sigma^2 J(e, e)$. Since $J(e, e) = 0$, we have $\alpha(U) = 0$. The latter part is as follows. $-\alpha(X)U + \nabla_X(U) = \frac{X(\sigma^{-1})}{\sigma^{-1}}U + \nabla_X(U) = \sigma\nabla_X(\sigma^{-1}U) = -c\,\sigma\nabla_X(e) = 0$. Thus we have $B(X) = 0$. Thereby $R^\sigma(X, Y) = 0$.

For a property $\nabla^\sigma e = 0$, we use the equation

$$\nabla^\sigma_X(e) - \nabla_X(e) = \alpha(X)e + (X \wedge U)(e) = J(U, X)e + J(U, e)X - J(X, e)U. \tag{5.45}$$

First we have $\nabla_X(e) = 0$. Since $U = -c\,\sigma\,e$, we have $J(U,X)e - J(X,e)U = 0$. Since $J(U,e) = -c\,\sigma(e,e) = 0$, we have $\nabla^{\sigma} e = 0$.

For a property of homogeneity conditions $Lie_E(J_a) = J_a$, we obtain it because $Lie_E(J) = J$ and $Lie_E(\sigma) = 0$.

We give a proof of "only if" part of (1). We assume that $(M, \circ, e, E, J_\sigma)$ is a Frobenius manifold.

We first show that $U = fe$ for some $f \in \Gamma(M, \mathcal{O}_M)$. By the conditions $\nabla_X e = \nabla^{\sigma}_X e = 0$ and (5.45), we have

$$J(U,X)e + J(U,e)X - J(X,e)U = 0. \tag{5.46}$$

We substitute $X = e$, we have $2J(U,e)e = 0$. Since $e \neq 0$, we have $J(U,e) = 0$. By (5.46), we have $J(U,X)e = J(X,e)U$. Since J is non-degenerate and flat, we take X such that $J(X,e)$ is non-zero constant. Then we have $U = fe$ for some $f \in \Gamma(M, \mathcal{O}_M)$.

We show that $U = -c\,\sigma\,e$ for some $c \in \mathbb{C}$. By (5.44), we have

$$B(X) \wedge Y + X \wedge B(Y) = 0. \tag{5.47}$$

We could easily check that if $\dim M \geq 3$ then (5.47) implies that $B(X) = 0$. Since $U = fe$, $\alpha(U) = J(U,U) = f^2 J(e,e) = 0$. Thus we have $-\alpha(X)U + \nabla_X(U) = 0$. Since $-\alpha(X)U + \nabla_X(U) = \sigma\nabla_X(\sigma^{-1}U) = \sigma\nabla_X(\sigma^{-1}fe) = \sigma X(\sigma^{-1}f)e$, we have $\sigma^{-1}f = -c$ for some $c \in \mathbb{C}$. Thus we have a result.

For a proof of (2), we see that $Lie_E(J(e)) = (D-1)J(e)$. Then we have $Lie_E(\sigma) = (1-D)\sigma$ and $Lie_E(\sigma^2 J) = (2 - 2D + D)(\sigma^2 J) = (2-D)(\sigma^2 J)$. ☐

Corollary 5.12. *For a conformal Frobenius manifold $(\circ, e, E, J)$ of a $\mathbb{C}^*$-bundle $p : L \to M$, we assume that $\iota(f)$ is a good section for a $\mathbb{C}^*$-equivariant morphism $f : L \to \mathbb{C}^*$. For $g : L \to \mathbb{C}^*$: a $\mathbb{C}^*$-equivariant morphism, $\iota(g)$ is a good section if and only if*

$$d((f/g)^{-1}) = c(\iota(f)^* J)(\iota(f)^* e) \tag{5.48}$$

for some $c \in \mathbb{C}$.

Proof. Since $\iota(g)^* J = (f/g)^2 \iota(f)^* J$ by Lemma 4.6, we obtain the result. ☐

5.11 Proof of Theorem 4.9

We may assume that $(\circ, e, E, J)$ be a conformal Frobenius structure constructed in §5.9. Then it has a good section $\iota(a)$. Let $h : D^2//W^2 \to \mathbb{C}^*$ be a $\mathbb{C}^*$-equivariant mapping whose section $\iota(h)$ is a good section of $(\circ', e', E, J')$.

Lemma 5.13. *By the embedding $(\mathrm{rad} I_F)_{\mathbb{C}} \hookrightarrow \Gamma(D^2//W^2, \mathcal{O}_{D^2//W^2})$ in Proposition 3.4(4), h is the image of this embedding.*

Proof. On $\mathbb{P}(\overset{\circ}{D}{}^2)//W^2$, $\iota(h)^* I^*_{D^2//W^2}$ and $\iota(a)^* I^*_{D^2//W^2}$ are non-degenerate dual metric. Also they are flat because the intersection form of the Frobenius manifold is flat ([3, p.191]).

Since

$$\iota(h)^* I^*_{D^2//W^2} = (a/h)^{-2} \iota(a)^* I^*_{D^2//W^2} \tag{5.49}$$

by Lemma 4.6, we apply Proposition 5.10 and we see (h/a) must be constant or suitable quardic or a suitable affine function with respect to flat coordinates for $\iota(a)^* I^*_{D^2//W^2}$. Also h/a must be an element of $\boldsymbol{\mathcal{S}}^W_{D^2,0,*}$ because the tensors $\iota(h)^* I^*_{D^2//W^2}$ and $\iota(a)^* I^*_{D^2//W^2}$ are homogeneous.

By the isomorphism $\mathbb{P}(D^2)//W^2 \simeq D^1_a//W^2$ of the diagram (5.30), $\iota(a)^* I^*_{D^2//W^2}$ corresponds to $I^*_{D^1//W^2}$ on $D^1//W^2$. Since flat coordinates for $I^*_{D^1//W^2}$ is an affine coordinate of D^1, we see that h/a must be of the form $x(b/a)a + y$ for $b \in (\mathrm{rad} I_F)_{\mathbb{C}} \setminus \mathbb{C}a$ and $x, y \in \mathbb{C}$. Thus we have a result. □

We remind the fact. For a Frobenius manifold $(M, \circ, e, E, J)$ with intersection form I^*, we put

$$\mathcal{F}_M(v) := \{\omega \in \Omega^1_M \mid Lie_v \omega = 0\} \tag{5.50}$$

for $v \in \Gamma(M, \Theta_M)$. In [13, Proposition 5.1], we have shown the fact that

$$e^2 I^*(\omega_1, \omega_2) = 0, \tag{5.51}$$

$$e I^*(\omega_1, \omega_2) = J^*(\omega_1, \omega_2) \tag{5.52}$$

for $\omega_1, \omega_2 \in \mathcal{F}_M(e)$. We have a following Lemma.

Lemma 5.14. *For a conformal Frobenius structure* $(\circ', e', E, J')$*, we have*

$$(\iota(a)^* e')^2 [\iota(a)^* I^*_{D^2//W^2}(\omega_1, \omega_2)] = 0 \tag{5.53}$$

for $\omega_1, \omega_2 \in \mathcal{F}_{\mathbb{P}(D^2)//W^2}(\iota(a)^* e')$.

Proof. Since $(\mathbb{P}(D^2)//W^2, \iota(h)^*\circ', \iota(h)^* e', \iota(h)^* E, \iota(h)^* J')$ is a Frobenius manifold with intersection form $\iota(h)^* I^*_{D^2//W^2}$, we have

$$(\iota(h)^* e')^2 [\iota(h)^* I^*_{D^2//W^2}(\eta_1, \eta_2)] = 0 \tag{5.54}$$

for $\eta_1, \eta_2 \in \mathcal{F}_{\mathbb{P}(D^2)//W^2}(\iota(h)^* e')$ by (5.51).

Since $\iota(a)^* e' = \iota(h)^* e'$ and $\iota(a)^* I^*_{D^2//W^2} = (h/a)^{-2} \iota(h)^* I^*_{D^2//W^2}$ by Lemma 4.6, we have

$$\begin{aligned}
&(\iota(a)^* e')^2 [\iota(a)^* I^*_{D^2//W^2}(\omega_1, \omega_2)] \\
=&(\iota(h)^* e')^2 [(h/a)^{-2} \iota(h)^* I^*_{D^2//W^2}(\omega_1, \omega_2)] \\
=&(h/a)^{-2} (\iota(h)^* e')^2 [\iota(h)^* I^*_{D^2//W^2}(\omega_1, \omega_2)] \\
=&0
\end{aligned}$$

for $\omega_1, \omega_2 \in \mathcal{F}_{\mathbb{P}(D^2)//W^2}(\iota(a)^* e')$, where we used the fact that $(\iota(h)^* e')(h/a) = 0$ because its degree is negative with respect to the Euler field $\iota(h)^* E$ by Lemma 5.13. □

Lemma 5.15. *There exists $c \in \mathbb{C}^*$ such that $e' = ce$, $J' = c^{-1}J$.*

Proof. Since $(\mathbb{P}(D^2)//W^2, \iota(a)^*\circ, \iota(a)^*e, \iota(a)^*E, \iota(a)^*J)$ is a Frobenius manifold with intersection form $\iota(a)^*I^*_{D^2//W^2}$, we have

$$(\iota(a)^*e)^2[\iota(a)^*I^*_{D^2//W^2}(\omega_1,\omega_2)] = 0 \tag{5.55}$$

for $\omega_1, \omega_2 \in \mathcal{F}_{\mathbb{P}(D^2)//W^2}(\iota(a)^*e)$ by (5.51).

As we have seen in [11], a non-singular vector field $v \in D^1//W^2$ of degree -1 satisfying

$$v^2 I^*_{D^1_a//W^2}(\omega_1,\omega_2) = 0 \tag{5.56}$$

for $\omega_1, \omega_2 \in \mathcal{F}_{D^1_a//W^2}(v)$ is unique up to $\mathbb{C}^*$-multiplication, see also [13, (4.13)]. By Lemma 5.14 and (5.55), we have $\iota(a)^*e' = c\iota(a)^*e$ for some $c \in \mathbb{C}^*$.

By (5.52), we have

$$(\iota(h)^*e')(\iota(h)^*I^*_{D^2//W^2})(\omega_1,\omega_2) = (\iota(h)^*J')^*(\omega_1,\omega_2) \text{ for } \omega_1,\omega_2 \in \mathcal{F}_{\mathbb{P}(D^2)//W^2}(\iota(h)^*e'),$$
$$(\iota(a)^*e)(\iota(a)^*I^*_{D^2//W^2})(\omega_1,\omega_2) = (\iota(a)^*J)^*(\omega_1,\omega_2) \text{ for } \omega_1,\omega_2 \in \mathcal{F}_{\mathbb{P}(D^2)//W^2}(\iota(a)^*e).$$

Since $\iota(h)^*e' = \iota(a)^*e' = c\iota(a)^*e$, we have $\mathcal{F}(\iota(h)^*e') = \mathcal{F}(\iota(a)^*e)$. Then we have

$$\begin{aligned}
&(\iota(a)^*J)^*(\omega_1,\omega_2)\\
=&(\iota(a)^*e)(\iota(a)^*I^*_{D^2//W^2})(\omega_1,\omega_2)\\
=&c^{-1}(\iota(h)^*e')[(h^2/a^2)\iota(h)^*I^*_{D^2//W^2}](\omega_1,\omega_2)\\
=&c^{-1}(h^2/a^2)(\iota(h)^*J')^*(\omega_1,\omega_2)\\
=&c^{-1}(\iota(a)^*J')^*(\omega_1,\omega_2)
\end{aligned}$$

for $\omega_1, \omega_2 \in \mathcal{F}(\iota(a)^*e)$. Since $\mathcal{F}(\iota(a)^*e)$ is sufficient to determine the tensor ([11]), we have $(\iota(a)^*J)^* = c^{-1}(\iota(a)^*J')^*$. By the correspondence of Lemma 4.6, we have $e' = ce$ and $J' = c^{-1}J$. □

Lemma 5.16. *$\iota(a)$ is a good section of $(\circ', e', E, J')$.*

Proof. By Lemma 5.13, $h \in (\mathrm{rad}I_F)_{\mathbb{C}}$. By Theorem 5.9, $d(b/a) = const.(\iota(a)^*J)(\iota(a)^*e)$. Thus we have $d(h/a) = c_1(\iota(a)^*J)(\iota(a)^*e)$ for some $c_1 \in \mathbb{C}$. Then we have

$$\begin{aligned}
d[(h/a)^{-1}] &= -(h/a)^{-2}d(h/a)\\
&= -(h/a)^{-2}[c_1(\iota(a)^*J)(\iota(a)^*e)]\\
&= -c_1(\iota(h)^*J)(\iota(h)^*e)\\
&= -c_1(\iota(h)^*J')(\iota(h)^*e') \quad (\because \text{ Lemma 5.15}).
\end{aligned}$$

Since $\iota(h)$ is a good section, we apply the Corollary 5.12, then we have a result. □

We give a proof of Theorem 4.9. Then $(\mathbb{P}(D^2)//W^2, \iota(a)^*\circ, \iota(a)^*e, \iota(a)^*E, \iota(a)^*J)$ and $(\mathbb{P}(D^2)//W^2, \iota(a)^*\circ', \iota(a)^*e', \iota(a)^*E, \iota(a)^*J')$ are Frobenius manifolds whose intersection forms are $\iota(a)^*I^*_{D^2//W^2}$. Since we proved in [13] that Frobenius manifold structure whose intersection form is $I^*_{D^1_a//W^2}$ and Euler field $E_{D^1_a//W^2}$ is unique up to $\mathbb{C}^*$ on $D^1_a//W^2$, we also have

$$(\iota(a)^*\circ, \iota(a)^*e, \iota(a)^*E, \iota(a)^*J) = (c^{-1}\iota(a)^*\circ', c\iota(a)^*e', \iota(a)^*E, c^{-1}\iota(a)^*J') \tag{5.57}$$

for some $c \in \mathbb{C}^*$. By the correspondence in Lemma 4.6, we obtain the result.

5.12 Proof of Proposition 4.10

By a uniqueness of a conformal Frobenius structure, we may assume that $(\circ, e, E, J)$ is a conformal Frobenius structure constructed in §5.9. Then $(\iota(a)^*J)(\iota(a)^*e)$ equals to $d(b/a)$ up to a constant multiple by Theorem 5.9. Since $\iota(a)$ is a good section, we apply Corollary 5.12 and we have a result.

References

[1] I. N. Bernšteĭn, O. V. Švarcman, *Chevalley's theorem for complex crystallographic Coxeter groups*, Funktsional. Anal. i Prilozhen. **12** (1978) no. 4, 79–80.

[2] I. N. Bernšteĭn, O. V. Švarcman, *Chevalley's theorem for complex crystallographic Coxeter groups and affine root systems*, Seminar on Supermanifolds 2, edited by Leites, 1986 No.22, Matem. Inst., Stockholoms Univ.

[3] B. Dubrovin, *Geometry of 2D topological field theories*, Integrable Systems and Quantum Groups (ed. by R. Donagi, et al.), Lecture Notes in Math. **1620**, Springer-Verlag (1996), 120–348.

[4] R. Friedman, J.W. Morgan, *Holomorphic principal bundles over elliptic curves. II. The parabolic construction*, J. Differential Geom. **56** (2000), no. 2, 301–379.

[5] C. Hertling, *Frobenius manifolds and moduli spaces for singularities*, Cambridge Tracts in Math. **151**, Cambridge Univ. Press, 2002.

[6] C. Houzel, *Géométrie analytique locale*, Séminaire Henri Cartan, 13ième année (1960/61).

[7] V. G. Kac, D. H. Peterson, *Infinite-dimensional Lie algebras, theta functions and modular forms*, Adv. in Math. **53** (1984) no. 2, 125–264.

[8] E. Looijenga, *Root systems and elliptic curves*, Invent. Math. **38** (1976/77) no.1, 17-32.

[9] K. Saito, *Period mapping associated to a primitive form*, Publ. RIMS, Kyoto Univ. **19** (1983) 1231–1264.

[10] K. Saito, *Extended affine root system I*, Publ. RIMS, Kyoto Univ. **21** (1985) 75-179.

[11] K. Saito, *Extended affine root system II*, Publ. RIMS, Kyoto Univ. **26** (1990) 15-78.

[12] I. Satake, *Automorphisms of the extended affine root system and modular property for the flat theta invariants*, Publ.RIMS Kyoto Univ. 31-1(1995), 1-32.

[13] I. Satake, *Frobenius manifolds for elliptic root systems*, arXiv:math/0611553.

[14] K. Wirthmuller, *Root systems and Jacobi forms*, Compositio Math. **82** (1992) no. 3, 293-354.